Generalitat de Catalunya

Programa de Prevenció y Atención a la Cronicidad de Cataluña

Qué hemos hecho y qué hemos aprendido

Sebastià J. Santaeugènia Gonzàlez, MD PhD

Programa de Prevenció i Atenció a la Cronicitat

Departament de Salut. Generalitat de Catalunya

Programa Interdepartamental d'Atenció i Interacció Social i Sanitària

Generalitat de Catalunya

1

Generalitat de Catalunya
**Programa de prevenció i atenció
a la cronicitat**

Aspecto nº1: Elaborando un nuevo entorno de salud

De qué va este nuevo entorno....

Las primeras cinco enfermedades causantes de muerte (75%) son crónicas y consumen más de la mitad de los recursos del sistema

Transición epidemiológica

Porcentaje de muerte por causa y sexo

Porcentaje de recursos económicos consumidos del sistema

El 53% de los recursos económicos del sistema son consumidos por las cinco principales causas de muerte, todas crónicas.

Envejecimiento poblacional

Ageing by numbers

What is the current situation for older people in your country? The coloured circles represent where 96 countries rank in the Global AgeWatch Index 2014 – the only global index to measure the social and economic wellbeing of older people

La cronicidad y la Edad

De las diferencias en el género y esperanza de vida

Age and Ageing Advance Access published August 4, 2013

Age and Ageing 2013; 0: 1–7
doi: 10.1093/ageing/aft115

© The Author 2013. Published by Oxford University Press on behalf of the British Geriatrics Society.
All rights reserved. For Permissions, please email: journals.permissions@oup.com

Cross-national disparities in sex differences in life expectancy with and without frailty

ROMAN ROMERO-ORTUNO¹, TONY FOUWEATHER², CAROL JAGGER²

Men die, women suffer?

Gender differences in health expectancy

REVES 2011,
Jan-Willem Bruggink,
Statistics Netherlands

Generalitat de Catalunya
Programa de prevenció i atenció
a la cronicitat

Health Women are sicker, men die quicker

Health Women are sicker, men die quicker

Women more illnesses – men die younger
In a nutshell, this is the main difference between women and men's health: men live an average of five years less than women, while women have more illnesses and health problems than men throughout their lives. This apparent paradox can be explained by a number of factors – both biological and social: it is assumed that men's lower mortality age is partly attributable to biological factors, but also to differences in lifestyle and various types of risk behaviour. The fact that women are more ill than men is often put down to differences in lifestyle and living conditions, for instance strains in relation to care work at home and in their professional lives.

Gender differences in life expectancy have lessened in recent years. Can this be an indication of a narrowing of the gap in health and mortality?

Fewer cardiovascular deaths – particularly among men
Cardiovascular diseases have long been the most common cause of death among men and women. In 2007, around 35 per cent of all deaths were caused by cardiovascular diseases, with a higher figure for men than women. However, a clear fall in cardiovascular deaths is emerging for both men and women, and the fall has been more pronounced for men. This decline is a contributing factor to the increase in life expectancy for both men and women.

With regard to the second most common cause of death – cancer – there have not been any major changes in the past 20 years, either for men or women. One in four deaths in 2007 was related to cancer. Cancer is one of the most common causes of death among the younger age groups in particular.

One in five smoke daily
The decline in cardiovascular diseases is assumed to be related to the decline in smoking: the proportion of men who smoke every day has fallen from more than 50 per cent at the start of the 1970s to 21 per cent in 2008. The corresponding figure for women was slightly higher than 30 per cent for a long time, but has now fallen to 21 per cent.

In Sweden, more women than men smoke on a daily basis – 17 and 12 per cent respectively in 2006. This is the trend in a number of countries among the younger age groups, with Norway, Sweden and Spain all having more women than men aged 15–24 smoking daily. It may therefore seem as if the numerous awareness campaigns in recent years have had an effect on middle-aged men, but have not prevented young girls from taking up smoking to the same extent.

Fewer dying in accidents
As is the case for cardiovascular diseases, the gender gap in accident-related deaths has also narrowed gradually. The number of fatalities due to accidents has fallen since 1980, both for men and women, with the biggest reduction for men.

Women more illnesses – men die younger
In a nutshell, this is the main difference between women and men's health: men live an average of five years less than women, while women have more illnesses and health problems than men throughout their lives. This apparent paradox can be explained by a number of factors – both biological and social: it is assumed that men's lower mortality age is partly attributable to biological factors, but also to differences in lifestyle and various types of risk behaviour. The fact that women are more ill than men is often put down to differences in lifestyle and living conditions, for instance strains in relation to care work at home and in their professional lives.

Gender differences in life expectancy have lessened in recent years. Can this be an indication of a narrowing of the gap in health and mortality?

Fewer cardiovascular deaths – particularly among men
Cardiovascular diseases have long been the most common cause of death among men and women. In 2007, around 35 per cent of all deaths were caused by cardiovascular diseases, with a higher figure for men than women. However, a clear fall in cardiovascular deaths is emerging for both men and women, and the fall has been more pronounced for men. This decline is a contributing factor to the increase in life expectancy for both men and women.

With regard to the second most common cause of death – cancer – there have not been any major changes in the past 20 years, either for men or women. One in four deaths in 2007 was related to cancer. Cancer is one of the most common causes of death among the younger age groups in particular.

One in five smoke daily
The decline in cardiovascular diseases is assumed to be related to the decline in smoking: the proportion of men who smoke every day has fallen from more than 50 per cent at the start of the 1970s to 21 per cent in 2008. The corresponding figure for women was slightly higher than 30 per cent for a long time, but has now fallen to 21 per cent.

In Sweden, more women than men smoke on a daily basis – 17 and 12 per cent respectively in 2006. This is the trend in a number of countries among the younger age groups, with Norway, Sweden and Spain all having more women than men aged 15–24 smoking daily. It may therefore seem as if the numerous awareness campaigns in recent years have had an effect on middle-aged men, but have not prevented young girls from taking up smoking to the same extent.

Fewer dying in accidents
As is the case for cardiovascular diseases, the gender gap in accident-related deaths has also narrowed gradually. The number of fatalities due to accidents has fallen since 1980, both for men and women, with the biggest reduction for men.

De Envejecer con desigualdad...

AGEING UNEQUALLY

Francesca Colombo
Head, OECD Health Division
Brussels, April 16

Inequality has increased in most OECD countries

Evolution of Gini coefficients between 1985 and 2010.

Source: OECD (2011), *Divided We Stand*

Being Old in 2040 Will Be No Fun

Current trends in demographics and national budgets will leave the elderly of the future in dire straits.

By Martin Hutchinson, March 31, 2015

Posts by Martin Hutchinson

Credit: John - www.flickr.com

There are times when it's good to be young – the 1960s, with its prosperity and hedonism, was one such period. Being old has fewer joys, but you can argue that the 1990s were a halcyon period for the old. Pension funds were swollen by stock appreciation and the senior citizen generation was relatively small.

However, with the current trends in demographics and national budgets, the younger generation of Baby Boomers and the older Gen-X'ers (only few of the older Baby Boomers will still be around) can rest assured of one thing: Being old in 2040 will be very unpleasant.

Growing gap in male life expectancy at age 65 by earnings levels in the U.S.

Source: Goldman and Orszag, 2014

Generalitat de Catalunya
Programa de prevenció i atenció
a la cronicitat

Dos reflexiones de futuro

JAMA JAMA
@JAMA_current

Age-adjusted prevalence of #obesity in 2013-2014 was 35.0% among men and 40.4% among women spr.ly/6011BSzwc

Survey Cycle	Sample Size	Odds Ratios (95% CI)		Predicted Margins, % (95% CI)	
		Adjusted for Age Group	Adjusted for Age Group, Race/Hispanic Origin Group, Smoking Status, and Educational Category	Adjusted for Age Group	Adjusted for Age Group, Race/Hispanic Origin Group, Smoking Status, and Educational Category
All^a					
2005-2006	4356	1 [Reference]	1 [Reference]	34.6 (31.9-37.4)	34.5 (31.7-37.4)
2007-2008	5550	0.97 (0.83-1.13)	0.96 (0.81-1.14)	33.9 (31.8-36.1)	33.6 (31.3-36.0)
2009-2010	5926	1.06 (0.92-1.22)	1.06 (0.91-1.23)	35.9 (34.1-37.7)	35.8 (34.1-37.5)
2011-2012	5181	1.02 (0.86-1.21)	1.04 (0.87-1.25)	35.1 (32.4-37.9)	35.4 (32.8-38.2)
2013-2014	5455	1.15 (1.00-1.33)	1.18 (1.01-1.37)	37.9 (36.2-39.7)	38.1 (36.3-40.0)
Total	26 468				
Men^d					
2005-2006	2237	1 [Reference]	1 [Reference]	33.5 (29.5-37.7)	33.4 (29.3-37.7)
2007-2008	2746	0.94 (0.75-1.18)	0.94 (0.74-1.20)	32.2 (29.4-35.1)	32.1 (29.1-35.2)
2009-2010	2889	1.10 (0.87-1.39)	1.09 (0.86-1.38)	35.5 (32.2-39.0)	35.3 (32.2-38.6)
2011-2012	2585	1.01 (0.81-1.25)	1.02 (0.82-1.28)	33.7 (31.1-36.4)	33.9 (31.4-36.6)
2013-2014	2638	1.08 (0.88-1.32)	1.08 (0.87-1.34)	35.2 (33.1-37.3)	35.1 (33.1-37.3)
Total	13 095				
Women^e					
2005-2006	2119	1 [Reference]	1 [Reference]	35.7 (33.0-38.5)	35.6 (33.0-38.3)
2007-2008	2804	0.99 (0.85-1.16)	0.97 (0.83-1.14)	35.5 (33.4-37.7)	35.0 (32.8-37.3)
2009-2010	3037	1.03 (0.89-1.18)	1.02 (0.88-1.18)	36.3 (34.5-38.1)	36.0 (34.2-37.8)
2011-2012	2596	1.04 (0.85-1.26)	1.06 (0.86-1.31)	36.5 (33.0-40.1)	36.9 (33.3-40.7)
2013-2014	2817	1.23 (1.04-1.45)	1.28 (1.08-1.51)	40.5 (37.9-43.2)	41.1 (38.5-43.7)
Total	13 373				

Abbreviation: BMI, body mass index.

^a BMI was calculated as weight in kilograms divided by height in meters squared. Obese was defined as participants with a BMI of 30 or greater.

^b All of the models for both sexes combined include sex as a covariate.

^c Models for both sexes, combined for all covariates, include significant interactions between race/Hispanic origin group and educational category.

status, race/Hispanic origin group and sex, educational category and sex, and smoking status and sex.

^d For men, the model includes an interaction between race/Hispanic origin group and smoking status.

^e For women, the model includes interactions between age group and educational category, race/Hispanic origin group and educational category, age group and smoking status, and race/Hispanic origin group.

SPECIAL REPORT

A Potential Decline in Life Expectancy in the United States in the 21st Century

S. Jay Olshansky, Ph.D., Douglas J. Passaro, M.D., Ronald C. Hershov, M.D., Jennifer Layden, M.P.H., Bruce A. Carnes, Ph.D., Jacob Brody, M.D., Leonard Hayflick, Ph.D., Robert N. Butler, M.D., David B. Allison, Ph.D., and David S. Ludwig, M.D., Ph.D.

VIEWPOINT

David S. Ludwig, MD, PhD
New Balance Foundation Obesity Prevention Center, Boston Children's Hospital and Harvard Medical School, Boston, Massachusetts.

Lifespan Weighed Down by Diet

Since the end of the Civil War until the late 20th century, lifespan increased rapidly in the United States, a tremendous public health triumph brought about by a more dependable food supply, improved sanitation, and advances in medical care. In 1850, life expectancy among whites was an estimated 38 years for men and 40 years for women. These numbers nearly doubled by 1980, to 71 years for men and 78 years for women. With the start of the obesity epidemic in the late 1970s, this trend began to slow, leading some to predict that life expectancy would decline in the United States by the mid-21st century.¹

those most severely affected by the obesity epidemic (ie, counties located predominately in the Southeast and Midwest). This downward trend in longevity will almost certainly accelerate as the current generation of children—with higher body weights from earlier in life than ever before—reaches adulthood. Modern medical care may prevent premature death among adults who develop obesity at age 45 years, diabetes at 55 years, and heart disease at 65 years, but the public health implications are likely vastly greater if this sequence of events were initiated in childhood.

In addition to the health-related effects, the eco-

Original Investigation

Trends in Obesity Among Adults in the United States, 2005 to 2014

Katherine M. Flegal, PhD; Deanna Kruszon-Moran, MS; Margaret D. Carroll, MSPH; Cheryl D. Fryar, MSPH; Cynthia L. Ogden, PhD

De determinantes sociales y enfermedad...

Personas que viven en zonas deprimidas adquieren enfermedades crónicas 10 años antes que viven en zonas menos deprimidas

MARCA MAS EL C.P. QUE LA CARGA DE MORBILIDAD

Hay **diferencias de esperanza de vida de hasta 11,3 años** entre **diferentes barrios ciudad de Barcelona**

Aspecto nº2: Creando El CAORDEN

De la MULTIMORBILIDAD a la CRONICIDAD, de la cronicidad a la COMPLEJIDAD

“CAORDEN”

Dícese del estado vital donde el Caos y el orden están en equilibrio.

Modelo Poblacional vs Grupo Población

■	Població de màxima complexitat (1%)
■	Població d'alt risc (4%)
■	Població de risc moderat (10%)
■	Població de baix risc (35%)
■	Població general (50%)

Modelo	Italia Vittorio Veneto	USA PACE	Canadá SIPA
Descripción General	Red de atención integrada	Hospital de día para anciano	Equipos MD base Comunitaria
Población Diana	Mayores 65 años Discapacidad con necesidad At Domicilio	Todo paciente anciano en domicilio con discapacidad	Ancianos frágiles que residen en Comunidad
Gestor Caso	SI	NO	SI
PTI	SI	SI	SI
Equipo Multidiscipli	SI	SI	SI
Valoración Holística	SI	SI	SI
Resultados	Menor Hospital. Mejor resultados salud	Menor Hospital. Menor Institucional.	Menor Hospital. Menor Institucional.

Santaeugenia S (2015). Tesis doctoral. Efectividad de un modelo de atención geriátrica integrado en el marco de una OSI. Universitat Autònoma de Barcelona, Barcelona.

La importancia de su impacto

% Pob.	%Mort.	%Ingres.	Gasto	% Gasto
1	28,9	138,4	12.273€	12,8%
4	8,4	33,5	5.326€	22,2%
10	1,3	7,5	2.321€	24,2%
35	0,2	2,6	850€	31,1%
50	0,0	0,4	188€	9,6%

EVOLUCIONAR DE ENFERMEDAD A MULTIMORBILIDAD ...

Global burden of COPD: risk factors, prevalence, and future trends

David M. Mannion, A. Conita Bickel

Diabetes Atlas

IDF Diabetes Atlas: Global estimates of the prevalence of diabetes for 2011 and 2030

David R. Whiting^{a,*}, Leonor Guariguata^a, Clara Weil^a, Jonathan Shaw^b

BMJ

BMJ 2013;346:f2510 doi: 10.1136/bmj.f2510 (Published 2 May 2013)

Page 1 of 4

ANALYSIS

Better management of patients with multimorbidity

Martin Roland and Charlotte Paddison call for greater emphasis on continuity of care and clinical judgment to improve the experience of patients with multiple conditions

Generalitat de Catalunya
Programa de prevenció i atenció
a la cronicitat

BMJ

BMJ 2012;345:e5205 doi: 10.1136/bmj.e5205 (Published 3 September 2012)

Page 1 of 10

RESEARCH

Managing patients with multimorbidity: systematic review of interventions in primary care and community settings

OPEN ACCESS

Susan M Smith *associate professor of general practice*¹, Hassan Soubhi *adjunct professor of family medicine*², Martin Fortin *professor of family medicine*², Catherine Hudon *associate professor of family medicine*², Tom O'Dowd *professor of general practice*³

Pla de Salut
de Catalunya 2016-2020

15

... Y AVANZAR HACIA LA ATENCION DE COMPLEJIDAD

BMJ

BMJ 2013;346:f2510 doi: 10.1136/bmj.f2510 (Published 2 May 2013)

Page 1 of 4

ANALYSIS

Better management of patients with multimorbidity

Martin Roland and Charlotte Paddison call for greater emphasis on continuity of care and clinical judgment to improve the experience of patients with multiple conditions

BMJ

BMJ 2012;345:e5205 doi: 10.1136/bmj.e5205 (Published 3 September 2012)

Page 1 of 10

RESEARCH

Managing patients with multimorbidity: systematic review of interventions in primary care and community settings

OPEN ACCESS

Susan M Smith associate professor of general practice¹, Hassan Soubhi adjunct professor of family medicine², Martin Fortin professor of family medicine³, Catherine Hudon associate professor of family medicine⁴, Tom O'Dowd professor of general practice⁵

NICE National Institute for Health and Care Excellence

NICE ya elabora guías y rutas asistenciales para personas con necesidades de atención sanitaria y social

Social care for older people with multiple long-term conditions overview

NICE Pathways bring together all NICE guidance, quality standards and other NICE information on a specific topic.

NICE Pathways are interactive and designed to be used online. They are updated regularly as new NICE guidance is published. To view the latest version of this pathway see:

<http://pathways.nice.org.uk/pathways/social-care-for-older-people-with-multiple-long-term-conditions>

Pathway last updated: 12 November 2015

<https://www.nice.org.uk/guidance/ng22>

The NEW ENGLAND JOURNAL of MEDICINE

HOME

ARTICLES & MULTIMEDIA

“HIGH-NEED, HIGH-COST PATIENTS”:

People with multiple chronic conditions that are often complicated by patients’ limited ability to care for themselves independently and by their complex social needs

Perspective

Caring for High-Need, High-Cost Patients — An Urgent Priority

David Blumenthal, M.D., M.P.P., Bruce Chernof, M.D., Terry Fulmer, Ph.D., R.N., John Lumpkin, M.D., M.P.H., and Jeffrey Selberg, M.H.A.

N Engl J Med 2016; 375:909-911 | September 8, 2016 | DOI: 10.1056/NEJMp1608511

Share:

Article

References

Citing Articles (1)

Metrics

Improving the performance of America's health system will require improving care for the patients who use it most: people with multiple chronic conditions that are often complicated by patients' limited ability to care for themselves independently and by their complex social needs. Focusing on this population makes sense for humanitarian, demographic, and financial reasons.

<http://bit.ly/2cPyLa7>

Audio Interview

A hablar de la complejidad... Intrínseca

A hablar de la complejidad... Extrínseca (Profesionales)

Kenneth Rockwood @Krockdoc · 11 h

To be expert in complexity is to accept uncertainty. In a complex system, any intervention has many consequences, not all knowable at start.

A hablar de la complejidad... Extrínseca (Sistema)

Plan de Crónicos de Catalunya: dos perfiles de complejidad

PCC:

Paciente con multimorbilidad o condición única que comporta gestión clínica difícil

2-3%

MACA:

Pronóstico de vida limitado, altas necesidades, orientación paliativa, planificación de decisiones avanzadas

1,5%

The screenshot shows the CatSalut patient portal interface. At the top, there are logos for Generalitat de Catalunya, Departament de Salut, and HC3. The patient's name, 'ROQUE RUIZ BUJANDA', and CIP number, 'RUBU0851023004', are displayed. Below this, there are sections for 'Dades pacient' (Patient Data) and 'Equip d'atenció primària' (Primary Care Team). The patient data section includes fields for name, CIP, date of birth, and document number. The primary care team section lists the primary care center and the attending physician. The interface is in Catalan and includes a search bar and navigation tabs.

Two patient cards are shown. The top card is labeled 'Pacient Crònic Complex' and the bottom card is labeled 'MACA'. Both cards display the patient's name, CIP number, and age (26 anys). The cards are part of a list of patients, with a search icon and a user profile icon visible on the right side of each card.

Paciente en situación de cronicidad compleja es identificado en base a **tres pasos clave**

Font: Programa de prevenció i atenció a la cronicitat. Departament de Salut. 2012.

Institut Català de la Salut **Pla d'Actuació Individualitzat** Data impressió: 04/04/2013
Darrera actualització: 04/04/2013

Data naixement: 04/06/1967 DNI: [REDACTED] NASS: [REDACTED]
Domicili [REDACTED]

Problemes de salut

Codi	Descripció	Data alta	Rellevància	Comentari
E78	TRASTORNS DEL METABOLISME DE LES LIPOPROTEINES I ALTRES LIPI	08/03/2013	Alta	
E11	DIABETIS MELLITUS TIPUS 2	05/03/2013	Alta	
F17.2	DEPENDÈNCIA PER CONSUM DE TABAC, SÍNDROME DE DEPENDÈNCIA	28/02/2013	Alta	Alta des d'Intel·ligència Activa
Z21	ESTAT D'INFECCIÓ ASIMPTOMÀTICA PEL VIRUS DE LA IMMUNODEFICIÈ	05/09/2012	Alta	
C16	TUMOR MALIGNE DE L'ESTÓMAC	27/07/2012	Alta	
B37.3	CANDIDIASI VAGINAL	19/01/2012	Alta	

Medicació activa

Inici	Medicament	Principi Actiu	Posologia	Durada
01/02/2013	ATROALDO 20MCG/PULSACION 10ML SOLUC INHALACION EN ENVASE A PRESION	IPRATROPI BROMUR	2 x 6 h.	Indefinida
01/02/2013	BEGLAN ACCUHALER 50MCG POLVO INHALACION 1 ACCUH + 1 TIRA 60 ALVEOLOS	SALMETEROL HIDROXINAFTOAT	1 x 12 h.	Indefinida
20/03/2013	IBUPROFENO ALDO-UNION 100MG/5ML 200ML SUSP ORA EFG	IBUPROFEN	1 x 24 h.	Indefinida
11/03/2013	SINTROM UNO GEIGY 1MG 60 COMPRIMIDOS	ACENOCUMAROL	20 x 1 s.	Indefinida
01/02/2013	TERBASMÍN TURBUHALER 0,5 MG/DOS 200 DOSIS POLV INH	TERBUTALINA	1 x 24 h.	Indefinida
07/02/2013	BISOLTUS 10MG/5ML 250ML SOLUCION	CODEINA CLORHIDRAT	1 x 24 h.	250 dies

Al·lèrgies

PS	Descripció	Qualificador	Reacció	Agent	Comentari
Z88.3	ALTRES ANTIINFECCIOSOS		Lieu		
Z88.0	LA PENICIL·LINA		Lieu		
Z88.7	SÉRUM O VACUNA		Important	vacunes grip	proves Esther

Directrius si crisi o descompensació **Pla de decisions anticipades**

Proves directrius Proves PDA

Atenció al pacient

· Servei teleassistència? Si No

· Atès per gestor de casos? Si No

· Viu en residència? Si No

· Atès per PADES? Si No

· Atès per ATDOM? Si No

· Viu sol? Si No

El PIIC como plan individualizado, interdisciplinario, compartido con la persona y disponible online para todo el sistema de salud

¿QUÉ TIENE?

¿QUÉ TOMA?

¿QUÉ HAY QUE HACER SI ...?

QUÉ PREFIERE

5 NIVELES DE ACTIVACIÓN:

Teléfono, domicilio inmediato, domicilio diferido
Atención intermedia, hospital de agudos

Aspecto nº4: REORGANIZANDO EL SISTEMA

© Organización Mundial de la Salud, 30-09-2015

La respuesta actual de salud pública: más de lo mismo no es suficiente.

- Se necesita **un cambio sistémico**. Los sistemas de salud suelen estar mejor **diseñados para curar enfermedades agudas** que para gestionar y reducir al mínimo las consecuencias de los estados crónicos prevalentes en la vejez .
- Además, en estos sistemas frecuentemente se **trabaja de manera compartimentada**, por lo que se abordan cada una de estas cuestiones por separado.
- Se suelen utilizar los **costosos servicios de atención de casos agudos para atender cuadros crónicos** y no se logra fomentar plenamente el funcionamiento de las personas mayores que reciben atención a largo plazo.

Y sólo hay una opción basada en la evidencia

GOV.UK Integrating health and social care in Torbay
Improving care for Mrs Smith

Local Government Association Rewiring Public Services
Integrated care value case toolkit

The Scottish Government Riaghaltas na h-Alba
Public service reform hub

Health and social care integration: do we make it work?
International best practice suggests successful, patient-integrated health and care systems share 11 common features

Guardian Professional, Wednesday 18 September 2013 10.00 BST
ehi ehealth INSIDER

The Belfast Birth of PARIS
The sharing of patient and client information between health and social care professionals, and creation of integrated care records are key features of the National Programme for IT (NPTI) in the English NHS, that the UK NHS has long had the aim of promoting the development of integrated systems to join up health and social care in South and East of England.

Co-ordinated care for people with complex chronic conditions: Key lessons and markers for success

The King's Fund and the Nuffield Trust
Developing a National Strategy for the Promotion of Integrated Care

The Evidence Base for Integrated Care

Nick Goodwin and Judith Smith

TheKingsFund Ideas that change health care

nuffieldtrust evidence for better health care

Sources: Kodner D, Goodwin N, Thiel V, King's Fund, 2011
Sources: Kodner D, Goodwin N, Thiel V, King's Fund, 2013

Crear sistemas de atención integrada social y sanitaria

An integrated care approach because...

Does Integrated Care Work?

Evidence from PACE, Torbay, PRISMA and Rovereto—and other programmes and pilots in North America, UK, Europe, and Australia specifically targeted to frail older people—strongly suggests that integrated care is capable of achieving positive outcomes, although it is not always clear which combination of strategies—and under what circumstances—produce the best results:

- Expanded service access, including primary care +++
- Enhanced co-ordination and continuity +++
- Improved health and functional status ++
- Reduced hospitalisation/nursing home admission/LOS ++
- Improved user experience, quality of life (QoL), and satisfaction +++
- Reduced carer burden ++
- Greater efficiency +
- Controlled/reduced costs +
- Perceived improvements in partnership working; also greater focus on governance and guidelines. +++

(Kodner, 2011)

Source: Kodner D. King's Fund, 2011

<http://www.kingsfund.org.uk/sites/files/kf/dennis-kodner-integrated-care-frail-older-people-kingsfund-sept12.pdf>

...con elementos concretos para facilitar la integración

Font: Programa de prevenció i atenció a la cronicitat. Departament de Salut. 2013.

...con elementos concretos para facilitar la integración (II)

Font: Programa de prevenció i atenció a la cronicitat. Departament de Salut. 2013.

Aspecto 4: La Historia clínica compartida como herramienta clave de comunicación

THIS IS THE **ELECTRONIC MEDICAL RECORD** THAT COLLECTS ALL IMPORTANT INFORMATION ON THE STATE AND EVOLUTION OF A PATIENT THROUGHOUT THEIR HEALTHCARE PROCESS

Sistema catalán es complejo...

- **Modelo multiproveedor** (> 160 proveedores – 32 IT systems) integrado en una sola red pública
- **Permite la gestión autónoma** de cada proveedor
- **Los proveedores son libres de seleccionar sus sistemas de información**
- **El 94% de los centros de atención primaria tienen el mismo sistema (eCAP)**
- **La interoperabilidad debe garantizarse**

Estadístiques /
Processos Batch

Portales
documentación HC3

Plataforma imagen digital
(repositorio de imágenes)

HCDSNS
Historia Clínica Digital
Sistema Nacional de Salud

epSOS
Smart Open Services
for European Patients

Centro
Sanitario

Profesionales

Administración

Ciudadano

Historia Clínica Compartida de Cataluña

Generalitat de Catalunya
Departament de Salut

CatSalut

HC3 Historia Clínica Compartida a Catalunya

Resum HC

Alertes actives AP		Problemes Actives AP		Condicionants de salut AP	
Data	Diagnòstic	Data	Diagnòstic	Data	Diagnòstic
29-04-2013	Antecedents personals d'al·lèrgia a la penicil·lina	01-01-2012	Consum de tabac	01-01-2012	Antecedents familiars de Trastorns mentals i tract
03-06-2012	Informe d'infermeria a l'alta d'internament	29-04-2013	Contingut	01-09-2011	Trastorns mentals i tract
22-05-2013	Informe mèdic alta internament	12-04-2013	Immunitzacions	01-08-2011	Trastorns mentals i tract
25-04-2013	Linfoma no Hodgkin: Cèl·lules mítics dividides petites i grans, folicular	01-04-2013	Alta atenció especialitzada ambulatòria	No s'han trobat dades	
12-03-2013	Anatomia patològica	14-05-2013	Densitometria òssia	Prescripció activa (Recepta electrònica)	
				17-12-2012	STIL BASS 310 TALLA 1 MEDIA
				17-12-2012	CORTA (A-B) COMP FURTE
				17-12-2012	SUBRO FISIOLÒGIC RIEF MEDICAL 50ML I FRAS
				17-12-2012	LOVAS TATENA CENTRUM 20MG 20 COMPRIMIDOS EFG

Catàlegs corporatius y aplicaciones Departamento de Salut

- Asegurados
- Centros sanitarios
- Auditoria
- CMBD Hospitalario
- Catàlegs diagnòstics
- Receta electrónica
- Registro del cáncer
- Voluntades anticipadas
- Seguridad y acceso

Repositorios externos centros sanitarios

- CENTRO N
- ...

Entidades certificadoras

- Catcert
- e-DNI

HC3 Deployment

SISCAT CENTRES WITH AN AGREEMENT WITH HC3

152
Millions of published documents

99%
centres SISCAT connected

665
Health Care Centres connected

Epidemiologists, ASP, ASPB and SEM

Other Departments in process: Social Welfare and Family

HC3 Publication

Aspecto 6: Inclusión de la complejidad en Salud Mental

Enfermedades crónicas:
30% de la población

Trastornos mentales:
20% de la población

PCC en SM y Adicciones: Perfiles de cronicidad

- **Perfil 1.** Concurrencia de dos o más patologías mentales identificadas con criterio 1.
- **Perfil 2.** Concurrencia de una patología de criterio 1 y como mínimo una de criterio 2.
- **Perfil 3.** Concurrencia de una patología del criterio 1 y condiciones sociales adversas.
- **Perfil 4.** Concurrencia de una patología criterio 1 y comorbilidad física relevante
- **Perfil 5.** Personas desvinculadas de APS y de SM.
 - Personas con trastorno mental crónico sin contacto en 1 año
 - Personas atendidas más de 10 veces en un servicio de UCIAS, en los últimos 6 meses.
- **Perfil 6.** Cualquier condición de SM de difícil manejo, a criterio profesional

Clasificación de las personas atendida en los servicios de SM, según cronicidad y complejidad

No se incluyen adicciones

Resultados nº1: ¿COMO se COMPORTAN?

PCC

Paciente con multimorbilidad o condición única que comporta gestión clínica difícil

2-3%

MACA

Pronóstico de vida limitado, altas necesidades, orientación paliativa, planificación de decisiones avanzadas

1,5%

N= 98.000 personas	No PCC	PCC	MACA
Tasa mortalidad (x100)	0,8	12,8	42,4
Media visitas AP	5,8	22,9	20,7
Usuario Trabajo social sanitario (%)	2,8	35,4	30,8
Proporción en at. Domiciliaria (%)	3,9	48,2	63,6
Proporción usuarios servicios sociales (%)	0,9	13,4	30,8
Tasa ingresos urgentes (x100)	4,9	41,0	52,3
Proporción pacientes reingresadores (%)	0,7	8,9	11,4
Media visitas a CCEE	1,5	4,3	3,1
Media fármacos	3,7	12,6	11,2

Resultados nº1: ¿COMO se COMPORTAN? (II)

Población Catalunya

PCC

MACA

Resultados nº2: Calidad o Cantidad

Prevalencia personas en situación complejidad

■ MACA ■ PCC ■ TAXA AGREGADA

Resultados nº3: Aprendemos y por ello mejoramos

Prevalencia personas con PIIC bien hecho

Resultados nº3: Aprendemos y por ello mejoramos

Ayudando a Atención Primaria desde Geriátría, Cuidados Paliativos y Salud Mental

Rutas Asistenciales - Presente y Futuro

- ❑ Reflexión estado actual y Detección de los **déficits**
- ❑ Inclusión **ámbito residencial y Salud Mental**
- ❑ Generación del **modelo RAC 2.0 (+ operativo!)**
- ❑ Mejorar la capacidad de evaluación:
 - Evaluación accesible y transparente
 - Detección de buenas prácticas

territoriales

Ruta assistencial
Diabetis mellitus tipus 2

SAP Barcelonès Nord Maresme, ICS
Hospital Universitari Germans Trias i Pujol, ICS
Badalona Serveis Assistencials, BSA

Resultados nº4: Rutas Asistenciales

Taula 72. Resum dels ítems usats per avaluar el disseny de la RAC per a cada agrupació territorial amb RAC disponible

Regió Sanitària	Agrupacions territorials RAC (n=42)	RAC amb		Participació en la redacció de la RAC					
		PCC ¹	MACA ²	AP	HA	061	RES	CSS	SSMN
	Garraf								
	Vallès Oriental Central								
	Baix Vallès								
	Baix Montseny								
	Maresme Central								
Barcelonès Nord i Baix Maresme	Badalona ICS								
	Badalona BSA								
	Sta. Coloma								
	Baix Llob Litoral-Sant Boi								
	Baix Llob Centre-LHO nord								
	Baix Llob Litoral-Viladecans								
	El Prat i Hospitalet Sud								
	Baix Llobregat Nord								
	Vallès Occidental Est								
	Vallès Occidental Oest								
	Barcelona Dreta								
	Barcelona Esquerra								
	Barcelona Litoral Mar								
	Barcelona Nord								
36 AT amb RAC	Cobertura poblacional	96,4%	93,9%	85,1%	85,1%	52,4%	47,4%	79,4%	13,9%

Resultados nº4: Rutas Asistenciales

Descripció del grup	Núm. de pacients
Pacients crònics complexos	528 pacients
Malalts crònics avançats	134 pacients
Insuficiència	22 pacients
Ruta diabetis	11128 pacients
Ruta depressió	4771 pacients
Ruta Insuficiència Cardíaca	1262 pacients
Ruta MPOC	2566 pacients

Procés

Ruta diabetis Procés

(10.293 pacients)	indicador 1-a	indicador 1-b	Total Ind 1	%	2	%	3	%	4	%	indicador FU	%	sense HbA1c	%	
EAP Lluçanès	347	104	157	261	75,2%	256	73,8%	269	77,5%	179	51,6%	122	35,2%	86	24,8%
EAP Centelles	688	208	276	484	70,3%	452	65,7%	495	71,9%	216	31,4%	134	19,5%	204	29,7%
EAP La Vall del Ges - Torelló	1258	463	424	887	70,5%	965	76,7%	998	79,3%	736	58,5%	350	27,8%	371	29,5%
EAP Manlleu	1506	471	528	999	66,3%	1110	73,7%	1153	76,6%	744	49,4%	440	29,2%	507	33,7%
EAP Roda de Ter	791	264	267	531	67,1%	588	74,3%	606	76,6%	381	48,2%	206	26,0%	260	32,9%
EAP Sant Hipòlit de Voltregà	436	140	165	305	70,0%	325	74,5%	347	79,6%	191	43,8%	84	19,3%	131	30,0%
EAP Sant Quirze de Besora	441	143	126	269	61,0%	309	70,1%	334	75,7%	193	43,8%	102	23,1%	172	39,0%
EAP Santa Eugènia de Berga	1077	447	291	738	68,5%	822	76,3%	869	80,7%	593	55,1%	273	25,3%	339	31,5%
EAP Tona	770	291	230	521	67,7%	555	72,1%	593	77,0%	352	45,7%	208	27,0%	249	32,3%
EAP Vic 1 - Nord	1242	443	505	948	76,3%	959	77,2%	1029	82,9%	593	47,7%	444	35,7%	294	23,7%
EAP Vic 2 - Sud	1233	320	674	994	80,6%	963	78,1%	1014	82,2%	793	64,3%	282	22,9%	239	19,4%
Sense especificar	14	0	3	3	21,4%	3	21,4%	4	28,6%	2	14,3%	1	7,1%	11	78,6%
10.292	3.305	3.672	6.977	67,8%	7.347	71,4%	7.760	75,4%	4.993	48,5%	2.651	25,8%	3315	32,2%	

Els indicadors són els següents:

Indicador 1-a: Pacients amb diagnòstic DM2 i 2 determinacions HbA1c en un any

Indicador 1-b: Pacients amb diagnòstic DM2 i 1 determinacions HbA1c en un any

Indicador 2: Pacients amb diagnòstic DM2 i 1 determinació LDL-c en un any

Indicador 3: Pacients amb diagnòstic DM2 i que tenen resultat laboratori FG

Indicador 4: Pacients amb diagnòstic DM2 i que tenen quocient Alb/Creatina en orina

Indicador FU: Pacients amb Fons Ull

Tasa Hospitalización potencialmente evitable en complicaciones EPOC

Disminución 13,8% entre Dic. 2011 a Dic. 2014 (36 meses)

x 100.000 hab.

Fuente: MSIQ, Catsalut

Tasa Hospitalización potencialmente evitable en Ins Cardíaca

Disminución 2% entre Dic. 2011 a Dic. 2014 (36 meses)

Nueva tendencia!
Incremento del 27% desde 2006 a 2011

x 100.000 hab.

Fuente: MSIQ, Catsalut

Reingresos a 30 días para un conjunto procesos crónicos

Disminución 8,5% entre Dic. 2011 a Dic. 2015 (4 años)

Incluye: EPOC, Ins. Cardíaca, DM, asma, enf. coronarias

Generalitat de Catalunya
Departament de Salut

MSIQ: <http://146.219.25.61/msiq/index.html>

Pla de Salut
de Catalunya 2016-2020

Hospitalizaciones enf. crónicas en La Garrotxa

Comparativa con el resto de AGA (áreas)
1ª posición en 2015

“La Garrotxa” tiene nivel muy alto desarrollo servicios sociales (“Torbay catalán”)

Ventaja nº6: LA COMPLEJIDAD FUENTE DE INNOVACIÓN

Los Profesionales como agentes innovadores

Identificació d'una experiència de valor

Experiència de gestió o clínica que pot tenir valor per al sistema sanitari català, extreta a partir de:

Selecció en base a l'impacte de l'experiència

Experiència amb un impacte elevat en els proveïdors en què ha estat implementada

Selecció en base a la qualitat de l'avaluació i l'impacte

Experiència que ha realitzat una autoavaluació de qualitat i que ha tingut un impacte elevat en els proveïdors

Identificació d'elements d'èxit que configuren els projectes sistèmics

Experiència destacada

Experiència certificada

Projecte sistèmic

248
experiències registradas

48
Certificadas implantables

02/10/2014
Hospitalització Domiciliària Integral (HDI): experiència col·laborativa entre un Servei Hospitalari de Geriatria i un Servei de Rehabilitació Domiciliària
Sebastià Santaegúnia González

18/11/2016
Atenció a les persones amb necessitats complexes. Un model territorial a Osona
Isabel Ramon Bofarull

02/12/2016
Model integrat i territorial d'un farmacèutic oncològic dins d'una unitat d'onco-hematològica en un hospital comarcal
Gemma Basagaña

01/12/2016
PIC: Programa Itinerari Cuidadors. una experiència conjunta des de salut i serveis socials
Geles Martínez Meca

Líneas estratégicas...

Fuente: PPAC 2017

Estrategia de Formación

2012- 2016 Formación presencial /on-line sanitaria.

2016-2020 Formación on-line MOOC , transversal social y sanitaria.

...de futuro...

HC3 como herramienta comunicativa bidireccional "clave"

Sistemas de información interoperables social y sanitarios

Desarrollo y implementación de nuevos contenidos *La Meva Salut*

Desarrollo y implementación de modelos de atención no presencial:
e.Consulta, telemonitorización, teleasistencia...

Sistema de información claro, transversal de "gestión caliente"

Estratificación (GMA) como herramienta de soporte a evolucionar

Atención Integrada: elementos nucleares para el nuevo modelo

Modelo de **governanza territorial** social i sanitari

ACP: empoderam./ autocuidado

“Microsistemas”:

- Orientación comunitaria
- Rutas asistenciales integradas
- Trabajo interdisciplinar
- Buenas transiciones
- Atención 7/24
- Impulso atención domiciliaria
- Buena atención residencias

Valoración y Plan intervención “compartido” + Gestión caso

Liderazgo clínico y profesional

Marco evaluativo compartido:

Responsabilidad compartida en resultados

Alineación incentivos

ELEMENTOS FACILITADORES

Modelos **estratificación** que incorporen datos sociales

Gestión del cambio y culturas organizativas

Historia clínica y social compartida

Estrategia multipalanca: TODO al mismo tiempo

Generalitat de Catalunya

Muchas gracias

@sjsantaeugenia

cronicitat.salut@gencat.cat