



Informes Periódicos de Orphanet

serie Medicamentos Huérfanos

Octubre 2012

Listados de Medicamentos Huérfanos en Europa

Con designación huérfana en Europa y autorización de comercialización europea*

Con autorización de comercialización europea*, sin previa designación huérfana
en Europa

*Autorización de comercialización de la Comunidad Europea mediante el procedimiento centralizado

www.orphanet.es

Indice

Metodología	3
Listado de medicamentos huérfanos en Europa con designación huérfana europea y autorización de comercialización europea	4
1- Por nombre comercial en orden alfabético.....	4
2- Por fecha de la AC en orden descendiente.....	14
3- Por categoría ATC.....	15
4- Por el titular de la AC.....	16
Listado de medicamentos huérfanos en Europa con autorización de comercialización europea, sin previa designación huérfana en Europa	17
1- Por nombre comercial en orden alfabético.....	17
2- Por fecha de la AC en orden descendiente.....	29
3- Por categoría ATC.....	30
4- Por el titular de la AC.....	31

Metodología

Este documento proporciona un listado de todos los medicamentos huérfanos que, hasta la fecha indicada en el documento, han recibido una autorización de comercialización (AC) europea. Estos productos sanitarios pueden ser a partir de ahora accesibles en algunos países europeos, aunque no necesariamente en todos. En realidad, la accesibilidad de un medicamento huérfano concreto en un determinado país depende de la estrategia del laboratorio y de la decisión tomada por parte de las autoridades sanitarias nacionales respecto al reembolso.

Listado de medicamentos huérfanos en Europa con designación huérfana europea y autorización de comercialización europea.

En sentido estricto, los medicamentos huérfanos en Europa son aquellos fármacos a los que se les ha concedido una designación huérfana europea (de acuerdo con la regulación (EC) No 141/2000), y a los que se les ha concedido también una autorización de comercialización europea y, eventualmente, una evaluación positiva con un beneficio significativo.

Este listado se obtiene cruzando el listado de productos sanitarios que han recibido una designación huérfana (<http://ec.europa.eu/health/documents/community-register/html/alforphreg.htm>) con el listado de productos sanitarios que han recibido una autorización de comercialización (<http://ec.europa.eu/health/documents/community-register/html/alfregister.htm>).

Ambos listados están disponibles en el sitio web de la Dirección General de Sanidad y Consumo (DG Sanco) de la Comisión Europea.

El listado de medicamentos está ordenado alfabéticamente por nombre comercial.

La información que se proporciona incluye el nombre comercial, el nombre de la sustancia activa, la indicación de la autorización de comercialización (AC), y el titular y la fecha de la AC.

Para permitir una búsqueda multi-criterio, se proponen tres listados adicionales, clasificados :

- por fecha de la AC en orden descendiente.
- por categoría ATC
- por el titular de la AC

Listado de medicamentos huérfanos en Europa con autorización de comercialización europea, sin previa designación huérfana en Europa.

En sentido amplio, la denominación "medicamentos huérfanos" se usa para productos sanitarios a los que se les ha concedido una autorización de comercialización europea, pero que no han recibido una designación huérfana europea o cuya designación ha sido retirada. A estos fármacos se les puede haber concedido, o no, una designación huérfana en otras áreas geográficas del mundo.

En todos los casos, se les ha concedido una autorización de comercialización europea para una o más indicaciones de uso para una enfermedad rara, y aparecen en el listado de fármacos de la Dirección General de Sanidad y Consumo de la Comisión Europea a los que se les ha concedido una autorización de comercialización : <http://ec.europa.eu/health/documents/community-register/html/alfregister.htm>

El listado de medicamentos está ordenado alfabéticamente por nombre comercial.

La información que se proporciona incluye el nombre comercial, el nombre de la sustancia activa, la enfermedad rara para la que la autorización de comercialización (AC) está indicada, y el titular y la fecha de la AC.

Para permitir una búsqueda multi-criterio, se proponen tres listados adicionales, clasificados :

- por fecha de la AC en orden descendiente.
- por categoría ATC
- por el titular de la AC

Existe información adicional de cada producto en la pestaña "Medicamentos huérfanos" del sitio web de Orphanet www.orphanet.es o en el sitio web de la Agencia Europea de Medicamentos (EMA) <http://www.ema.europa.eu>. El listado de la EMA cubre todos los productos sanitarios autorizados en el mercado, no únicamente los medicamentos huérfanos. Los medicamentos huérfanos con designación huérfana europea están indicados con el logo .

Para cualquier pregunta o comentario, por favor contacte con: contact.orphanet@inserm.fr

Listado de medicamentos huérfanos en Europa con designación huérfana europea y autorización de comercialización europea

1- Por nombre comercial en orden alfabético

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
AFINITOR	Everolimus	This product is no longer an orphan medicine. It was originally designated an orphan medicine on 5 June 2007. Upon request of the marketing authorisation holder, Afinitor has now been removed from the Community Register of orphan medicinal products. Cf "List of orphan drugs in Europe with European market authorisation without prior orphan designation in Europe"		
ALDURAZYME	Laronidase	Long-term enzyme replacement therapy in patients with a confirmed diagnosis of Mucopolysaccharidosis I (MPS I; a [alpha]-L-iduronidase deficiency) to treat the non-neurological manifestations of the disease	10/06/2003	Genzyme Europe B.V.
ARZERRA	Ofatumumab	Treatment of chronic lymphocytic leukaemia (CLL) in patients who are refractory to fludarabine and alemtuzumab	19/04/2010	Glaxo Group Ltd
ATRIANCE	Nelarabine	Treatment of patients with T-cell acute lymphoblastic leukaemia (T-ALL) and T-cell lymphoblastic lymphoma (T-LBL) whose disease has not responded to or has relapsed following treatment with at least two chemotherapy regimens	22/08/2007	Glaxo Group Ltd
BRONCHITOL	Mannitol	Bronchitol is indicated for the treatment of cystic fibrosis (CF) in adults aged 18 years and above as an add-on therapy to best standard of care	13/04/2012	Pharmaxis Pharmaceuticals Limited
BUSILVEX	Busulfan (Intravenous use)	Followed by cyclophosphamide (BuCy2) is indicated as conditioning treatment prior to conventional haematopoietic progenitor cell transplantation (HPCT) in adult patients when the combination is considered the best available option Followed by cyclophosphamide (BuCy4) or melphalan (BuMel) is indicated as conditioning treatment prior to conventional haematopoietic progenitor cell transplantation in paediatric patients	09/07/2003	Pierre Fabre Médicament
CARBAGLU	Carglumic acid	Treatment of hyperammonaemia due to N-acetylglutamate synthase primary deficiency , hyperammonaemia due to isovaleric acidaemia , hyperammonaemia due to methylmalonic acidaemia , hyperammonaemia due to propionic acidaemia	24/01/2003	Orphan Europe S.a.r.l.
CAYSTON	Aztreonam	Suppressive therapy of chronic pulmonary infections due to <i>Pseudomonas aeruginosa</i> in patients with cystic fibrosis (CF) aged 6 years and older	21/09/2009	Gilead Sciences International Limited
CEPLENE	Histamine dihydrochloride	Maintainance therapy for adult patients with acute myeloid leukaemia in first remission concomitantly treated with interleukin-2 (IL-2). The efficacy of Ceplene has not been fully demonstrated in patients older than age 60	07/10/2008	EpiCept GmbH

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
CYSTADANE	Betaine anhydrous	Adjunctive treatment of homocystinuria , involving deficiencies or defects in cystathionine beta-synthase (CBS), 5,10-methylene-tetrahydrofolate reductase (MTHFR), cobalamin cofactor metabolism (cbl). Cystadane should be used as supplement to other therapies such as vitamin B6 (pyridoxine), vitamin B12 (cobalamin), folate and a specific diet	15/02/2007	Orphan Europe S.a.r.l.
DACOGEN	Decitabine	Treatment of adult patients aged 65 years and above with newly diagnosed de novo or secondary acute myeloid leukaemia (AML), according to the World Health Organization (WHO) classification, who are not candidates for standard induction chemotherapy	20/09/2012	Janssen-Cilag International N V
DIACOMIT	Stiripentol	Use in conjunction with clobazam and valproate as adjunctive therapy of refractory generalized tonic-clonic seizures in patients with severe myoclonic epilepsy in infancy (SMEI, Dravet's syndrome) whose seizures are not adequately controlled with clobazam and valproate	04/01/2007	Biocodex
ELAPRASE	Idursulfase	Long-term treatment of patients with Hunter syndrome (Mucopolysaccharidosis II, MPS II)	08/01/2007	Shire Human Genetic Therapies AB
ESBRIET	Pirfenidone	In adults for the treatment of mild to moderate Idiopathic Pulmonary Fibrosis (IPF)	28/02/2011	InterMune UK Ltd.
EVOLTRA	Clofarabine	Treatment of acute lymphoblastic leukaemia (ALL) in paediatric patients who have relapsed or are refractory after receiving at least two prior regimens and where there is no other treatment option anticipated to result in a durable response	29/05/2006	Genzyme Europe B.V.
EXJADE	Deferasirox	Treatment of chronic iron overload due to frequent blood transfusions (≥ 7 ml/kg/month of packed red blood cells) in patients with beta thalassaemia major aged 6 years and older. Treatment of chronic iron overload due to blood transfusions when deferoxamine therapy is contraindicated or inadequate in the following patient groups: - in patients with beta thalassaemia major with iron overload due to frequent blood transfusions in (≥ 7 ml/kg/month of packed red blood cells) patients aged 2 to 5 years - in patients with beta thalassaemia major with iron overload due to infrequent blood transfusions (< 7 ml/kg/month of packed red blood cells) aged 2 years and older, - in patients with other anaemias aged 2 years and older.	28/08/2006	Novartis Euro-pharm Ltd
FABRAZYME	Recombinant human alpha-galactosidase A INN = Agalsidase beta	Long-term enzyme replacement therapy in patients with a confirmed diagnosis of Fabry disease (alpha-galactosidase A deficiency) <i>This orphan designated product has completed its 10 years of "market exclusivity"</i>	03/08/2001	Genzyme Europe B.V.

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
FIRAZYR	Icatibant acetate INN = Icatibant	Symptomatic treatment of acute attacks of hereditary angioedema (HAE) in adults (with C1-esterase-inhibitor deficiency)	11/07/2008	Shire Orphan Therapies GmbH
FIRDAPSE (ex-ZENAS)	Amifampridine	Symptomatic treatment of Lambert-Eaton myasthenic syndrome (LEMS) in adults	23/12/2009	Biomarin Europe Ltd
GLIOLAN	5-aminolevulinic acid hydrochloride	In adult patients for visualisation of malignant tissue during surgery for malignant glioma (World Health Organization grade III and IV)	07/09/2007	Medac GmbH
GLIVEC	Imatinib mesilate	<p>Treatment of :</p> <ul style="list-style-type: none"> - adult and paediatric patients with newly diagnosed Philadelphia chromosome (bcr-abl) positive (Ph+) chronic myeloid leukaemia (CML) for whom bone marrow transplantation is not considered as the first line of treatment - adult and paediatric patients with Ph+ CML in chronic phase after failure of interferon-alpha therapy, or in accelerated phase or blast crisis - adult patients with newly diagnosed Philadelphia chromosome positive acute lymphoblastic leukaemia (Ph+ ALL) integrated with chemotherapy - adult patients with relapsed or refractory Ph+ ALL as monotherapy - adult patients with myelodysplastic/myeloproliferative diseases (MDS/MPD) associated with platelet-derived growth factor receptor (PDGFR) gene re-arrangements - adult patients with advanced hypereosinophilic syndrome (HES) and/or chronic eosinophilic leukaemia (CEL) with FIP1L1-PDGFRα rearrangement - adult patients with Kit (CD 117) positive unresectable and/or metastatic malignant gastrointestinal stromal tumours (GIST) - adjuvant treatment of adult patients who are at significant risk of relapse following resection of Kit (CD117)-positive GIST. Patients who have a low or very low risk of recurrence should not receive adjuvant treatment - adult patients with unresectable dermatofibrosarcoma protuberans (DFSP) and adult patients with recurrent and/or metastatic DFSP who are not eligible for surgery <p><i>This orphan designated product has completed its 10 years of "market exclusivity" for its indication in chronic myeloid leukemia.</i> <i>For other indications, the sponsor has requested the removal of orphan designation from Community Register on 16 April 2012.</i></p>	07/11/2001	Novartis Euro-pharm Ltd
GLYBERA	Alipogene tiparvovec	For adult patients diagnosed with familial lipoprotein lipase deficiency (LPLD) and suffering from severe or multiple pancreatitis attacks despite dietary fat restrictions. The diagnosis of LPLD has to be confirmed by genetic testing. The indication is restricted to patients with detectable levels of LPL protein	29/10/2012	uniQure bio-pharma B.V.

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
ILARIS	Canakinumab	This product is no longer an orphan medicine. It was originally designated an orphan medicine on 20 March 2007. Upon request of the marketing authorisation holder, Ilaris has now been removed from the Community Register of orphan medicinal products. Cf "List of orphan drugs in Europe with European market authorisation without prior orphan designation in Europe"		
INCRELEX	Mecasermin	Long-term treatment of growth failure in children and adolescents with severe primary insulin-like growth factor 1 deficiency (Primary IGFD)	03/08/2007	Ipsen Pharma
INOVELON	Rufinamide	Adjunctive therapy in the treatment of seizures associated with Lennox Gastaut syndrome in patients aged 4 years and older	16/01/2007	Eisai Ltd
JAKAVI	Ruxolitinib	Treatment of disease-related splenomegaly or symptoms in adult patients with primary myelofibrosis (also known as chronic idiopathic myelofibrosis), post-polycythaemia-vera myelofibrosis or post-essential-thrombocythaemia myelofibrosis .	23/08/2012	Novartis Euro-pharm Ltd
KALYDECO	Ivacaftor	Kalydeco is indicated for the treatment of cystic fibrosis (CF) in patients age 6 years and older who have a G551D mutation in the CFTR gene	23/07/2012	Vertex Pharmaceuticals (U.K.) Limited
KUVAN	Sapropterin dihydrochloride INN = Sapropterin	Treatment of hyperphenylalaninaemia (HPA) in adult and paediatric patients of 4 years of age and over with phenylketonuria (PKU) who have been shown to be responsive to such treatment Treatment of hyperphenylalaninaemia (HPA) in adult and paediatric patients with tetrahydrobiopterin (BH4) deficiency who have been shown to be responsive to such treatment	02/12/2008	Merck Serono Europe Ltd
LITAK	Cladribine (subcutaneous use)	Treatment of hairy cell leukaemia	14/04/2004	Lipomed GmbH
LYSODREN	Mitotane	Symptomatic treatment of advanced (unresectable, metastatic or relapsed) adrenal cortical carcinoma	28/04/2004	Laboratoire HRA Pharma
MEPACT	Mifamurtide	In children, adolescents and young adults for the treatment of high-grade resectable non-metastatic osteosarcoma after macroscopically complete surgical resection. It is used in combination with post-operative multi-agent chemotherapy	06/03/2009	IDM Pharma SAS
MOZOBIL	Plerixafor	In combination with G-CSF to enhance mobilisation of haematopoietic stem cells to the peripheral blood for collection and subsequent autologous transplantation in patients with lymphoma and multiple myeloma whose cells mobilise poorly	31/07/2009	Genzyme Europe B.V.
MYOZYME	Recombinant human acid alpha-glucosidase INN = Alglucosidase alpha	Long-term enzyme replacement therapy (ERT) in patients with a confirmed diagnosis of Pompe disease (acid α -glucosidase deficiency)	29/03/2006	Genzyme Europe B.V.

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
NAGLAZYME	N-acetylgalacto-samine 4-sulfatase INN = Galsulfase	Long-term enzyme replacement therapy in patients with a confirmed diagnosis of Mucopolysaccharidosis VI (MPS VI; N-acetylgalactosamine 4-sulfatase deficiency; Maroteaux-Lamy syndrome)	24/01/2006	BioMarin Europe Ltd
NEXAVAR	Sorafenib tosylate Sorafenib	Treatment of hepatocellular carcinoma Treatment of patients with advanced renal cell carcinoma who have failed prior interferon-alpha or interleukin-2 based therapy or are considered unsuitable for such therapy	19/07/2006	Bayer Pharma AG
NOVOTHIR-TEEN	Catrideracog	Long-term prophylactic treatment of bleeding in patients 6 years and above with congenital factor XIII A subunit deficiency	03/09/2012	Novo Nordisk A/S
NPLATE	Romiplostim	Adult chronic immune (idiopathic) thrombocytopenic purpura (ITP) in splenectomised patients who are refractory to other treatments (e.g. corticosteroids, immunoglobulins). Nplate may be considered as second line treatment for adult non-splenectomised patients where surgery is contra-indicated	04/02/2009	Amgen Europe B.V.
ONSENAL	Celecoxib	Reduction of the number of adenomatous intestinal polyps in familial adenomatous polyposis (FAP), as an adjunct to surgery and further endoscopic surveillance This medicine is now withdrawn from use in the European Union, more information on: www.ema.europa.eu	17/10/2003	Pfizer Ltd
ORFADIN	Nitisinone	Treatment of patients with confirmed diagnosis of hereditary tyrosinemia type 1 (HT-1) in combination with dietary restriction of tyrosine and phenylalanine	21/02/2005	Swedish Orphan Biovitrum International AB
PEDEA	Ibuprofen	Treatment of a haemodynamically significant patent ductus arteriosus in preterm newborn infants less than 34 weeks of gestational age	29/07/2004	Orphan Europe S.a.r.l.
PEYONA (ex-NYMUSA)	Caffeine citrate	Treatment of primary apnea of premature newborns	02/07/2009	Chiesi Farmaceutici SpA
PHOTOBARR	Porfimer sodium (for use with photodynamic therapy)	Ablation of high-grade dysplasia (HGD) in patients with Barrett's Oesophagus This medicine is now withdrawn from use in the European Union, more information on: www.ema.europa.eu	25/03/2004	Pinnacle Biologics B.V.
PLENADREN	Hydrocortisone	Treatment of adrenal insufficiency in adults.	03/11/2011	ViroPharma SPRL
PRIALT	Ziconotide (intraspinal use)	Treatment of severe, chronic pain in patients who require intrathecal (IT) analgesia	21/02/2005	Eisai Ltd
REPLAGAL	Agalsidase alfa	Long-term enzyme replacement therapy in patients with a confirmed diagnosis of Fabry Disease (alpha-galactosidase A deficiency) This orphan designated product has completed its 10 years of "market exclusivity"	03/08/2001	Shire Human Genetic Therapies AB

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
REVATIO	Sildenafil citrate INN = Sildenafil	Treatment of adult patients with pulmonary arterial hypertension classified as WHO functional class II and III, to improve exercise capacity. Efficacy has been shown in primary pulmonary hypertension and pulmonary hypertension associated with connective tissue disease. Treatment of paediatric patients aged 1 year to 17 years old with pulmonary arterial hypertension. Efficacy in terms of improvement of exercise capacity or pulmonary haemodynamics has been shown in primary pulmonary hypertension and pulmonary hypertension associated with congenital heart disease. Revatio solution for injection is for the treatment of adult patients with pulmonary arterial hypertension who are currently prescribed oral Revatio and who are temporarily unable to take oral therapy, but are otherwise clinically and haemodynamically stable. Revatio (oral) is indicated for treatment of adult patients with pulmonary arterial hypertension classified as WHO functional class II and III, to improve exercise capacity. Efficacy has been shown in primary pulmonary hypertension and pulmonary hypertension associated with connective tissue disease	28/10/2005	Pfizer Ltd
REVESTIVE	Teduglutide	Treatment of adult patients with short-bowel syndrome . Patients should be stable following a period of intestinal adaptation after surgery	30/08/2012	Nycomed Danmark ApS
REVLIMID	Lenalidomide	In combination with dexamethasone, treatment of multiple myeloma patients who have received at least one prior therapy	14/06/2007	Celgene Europe Ltd
REVOLADE	Eltrombopag	This product is no longer an orphan medicine. It was originally designated an orphan medicine on 3 August 2007. Upon request of the marketing authorisation holder, Revolade has now been removed from the Community Register of orphan medicinal products. Cf "List of orphan drugs in Europe with European market authorization without prior orphan designation in Europe"		
RILONACEPT REGENERON (ex-ARCALYST)	Rilonacept	Treatment of Cryopyrin-Associated Periodic Syndromes (CAPS) with severe symptoms, including Familial Cold Autoinflammatory Syndrome (FCAS) and Muckle-Wells Syndrome (MWS), in adults and children aged 12 years and older <i>This medicine is now withdrawn from use in the European Union, more information on: www.ema.europa.eu</i>	23/10/2009	Regeneron UK Limited
SAVENE	Dexrazoxane	Treatment of anthracycline extravasation	28/07/2006	SpePharm Holding BV
SIGNIFOR	Pasireotide	Treatment of adult patients with Cushing's disease for whom surgery is not an option or for whom surgery has failed	24/04/12	Novartis Euro-pharm Ltd
SIKLOS	Hydroxycarbamide	Prevention of recurrent painful vaso-occlusive crises including acute chest syndrome in paediatric and adult patients suffering from symptomatic Sickle Cell Syndrome	29/06/2007	Addmedica

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
SOLIRIS	Eculizumab	Treatment of patients with: - Paroxysmal nocturnal haemoglobinuria (PNH). Evidence of clinical benefit of Soliris in the treatment of patients with PNH is limited to patients with history of transfusions. - Atypical haemolytic uremic syndrome (aHUS).	20/06/2007	Alexion Europe SAS
SOMAVERT	Pegvisomant	Treatment of patients with acromegaly who have had an inadequate response to surgery and/or radiation therapy and in whom an appropriate medical treatment with somatostatin analogues did not normalize IGF-I concentrations or was not tolerated	13/11/2002	Pfizer Ltd
SPRYCEL	Dasatinib	Treatment of adult patients with: - newly diagnosed Philadelphia chromosome positive (Ph+) chronic myelogenous leukaemia (CML) in the chronic phase. - chronic, accelerated or blast phase CML with resistance or intolerance to prior therapy including imatinib mesilate. - Ph+ acute lymphoblastic leukaemia (ALL) and lymphoid blast CML with resistance or intolerance to prior therapy.	20/11/2006	Bristol-Myers Squibb Pharma EEIG
SUTENT	Sunitinib malate Sunitinib	This product is no longer an orphan medicine. This product was originally an orphan designated on 10 March 2005. Upon request by the MAH, Sutent has now been removed from the Community Register of Orphan Medicinal products. Cf "List of orphan drugs in Europe with European market authorisation without prior orphan designation in Europe"		
TASIGNA	Nilotinib	150 mg: Treatment of adult patients with newly diagnosed Philadelphia chromosome positive chronic myelogenous leukaemia (CML) in the chronic phase. 200 mg: Treatment of adult patients with: - newly diagnosed Philadelphia chromosome positive chronic myelogenous leukaemia (CML) in the chronic phase - chronic phase and accelerated phase Philadelphia chromosome positive chronic myelogenous leukaemia (CML) with resistance or intolerance to prior therapy including imatinib. Efficacy data in patients with CML in blast crisis are not available.	19/11/2007	Novartis Euro-pharm Ltd
TEPADINA	Thiotepa	In combination with other chemotherapy medicinal products: 1) with or without total body irradiation (TBI), as conditioning treatment prior to allogeneic or autologous haematopoietic progenitor cell transplantation (HPCT) in haematological diseases in adult and paediatric patients; 2) when high dose chemotherapy with HPCT support is appropriate for the treatment of solid tumours in adult and paediatric patients. It is proposed that Tepadina must be prescribed by physicians experienced in conditioning treatment prior to haematopoietic progenitor cell transplantation.	15/03/2010	Adienne S.r.l.


TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
THALIDOMIDE CELGENE	Thalidomide	In combination with melphalan and prednisone as first line treatment of patients with untreated multiple myeloma , aged >/= 65 years or ineligible for high dose chemotherapy	16/04/2008	Celgene Europe Ltd
THELIN	Sitaxentan sodium	Treatment of patients with pulmonary arterial hypertension classified as WHO functional class III, to improve exercise capacity. Efficacy has been shown in primary pulmonary hypertension and in pulmonary hypertension associated with connective tissue disease. <i>This medicine is now withdrawn from use in the European Union, more information on: www.ema.europa.eu</i>	10/08/2006	Pfizer Ltd
TOBI PODHALER	Tobramycin	Suppressive therapy of chronic pulmonary infection due to Pseudomonas aeruginosa in adults and children aged 6 years and older with cystic fibrosis	20/07/2011	Novartis Europharm Limited
TORISEL	Temsirolimus	First-line treatment of adult patients with advanced renal cell carcinoma (RCC) who have at least three of six prognostic risk factors Treatment of adult patients with relapsed and / or refractory mantle cell lymphoma (MCL)	19/11/2007	Pfizer Limited
TRACLEER	Bosentan mono-hydrate INN = Bosentan	Treatment of pulmonary arterial hypertension (PAH) to improve exercise capacity and symptoms in patients with WHO functional class III. Efficacy has been shown in: Primary (idiopathic and familial) PAH, PAH secondary to scleroderma without significant interstitial pulmonary disease, PAH associated with congenital systemic-to-pulmonary shunts and Eisenmenger's physiology. Some improvements have also been shown in patients with PAH WHO functional class II. To reduce the number of new digital ulcers in patients with systemic sclerosis and ongoing digital ulcer disease <i>This orphan designated product has completed its 10 years of "market exclusivity" for its indication in pulmonary arterial hypertension</i>	15/05/2002	Actelion Registration Ltd
TRISENOX	Arsenic Trioxide	Induction of remission and consolidation in adult patients with relapsed/refractory acute promyelocytic leukaemia (APL), characterised by the presence of the t(15;17) translocation and/or the presence of the Pro-Myelocytic Leukaemia/ Retinoic-Acid Receptor-alpha (PML/RAR-alpha) gene. Previous treatment should have included a retinoid and chemotherapy <i>This orphan designated product has completed its 10 years of "market exclusivity"</i>	05/03/2002	Cephalon Europe
VENTAVIS	Iloprost	Treatment of patients with primary pulmonary hypertension , classified as NYHA functional class III, to improve exercise capacity and symptoms	16/09/2003	Bayer Schering Pharma AG

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
VIDAZA	Azacitidine	Treatment of adult patients who are not eligible for haematopoietic stem cell transplantation with: -intermediate 2 and high risk myelodysplastic syndromes (MDS) according to the International Prognostic Scoring System (IPSS) -chronic myelomonocytic leukaemia (CMML) with 10-29 % marrow blasts without myeloproliferative disorder -acute myeloid leukaemia (AML) with 20-30 % blasts and multi-lineage dysplasia, according to World Health Organisation (WHO) classification	17/12/2008	Celgene Europe Ltd
VOLIBRIS	Ambrisentan	Treatment of patients with pulmonary arterial hypertension (PAH) classified as WHO functional class II and III, to improve exercise capacity. Efficacy has been shown in idiopathic PAH (IPAH) and in PAH associated with connective tissue disease	21/04/2008	Glaxo Group Ltd
VOTUBIA	Everolimus	Treatment of patients aged 3 years and older with subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis complex (TSC) who require therapeutic intervention but are not amenable to surgery. The evidence is based on analysis of change in SEGA volume. Further clinical benefit, such as improvement in disease-related symptoms, has not been demonstrated.	02/09/2011	Novartis Euro-pharm Ltd
VPRIV	Velaglucerase alfa	Long-term enzyme replacement therapy (ERT) in patients with type 1 Gaucher disease	26/08/2010	Shire Pharmaceuticals Ireland Ltd
VYNDAQEL	Tafamidis	Treatment of transthyretin amyloidosis in adult patients with stage 1 symptomatic polyneuropathy to delay peripheral neurologic impairment.	16/11/2011	Pfizer Specialty UK Ltd
WILZIN	Zinc acetate dihydrate	Treatment of Wilson's disease	13/10/2004	Orphan Europe S.a.r.l.
XAGRID	Anagrelide hydrochloride INN = Anagrelide	Reduction of elevated platelet counts in at risk essential thrombocythaemia patients who are intolerant to their current therapy or whose elevated platelet counts are not reduced to an acceptable level by their current therapy	16/11/2004	Shire Pharmaceutical Contracts Ltd
XALUPRINE (ex-MERCAP-TOPURINE NOVA)	Mercaptopurine	Indicated for the treatment of acute lymphoblastic leukaemia (ALL) in adults, adolescents and children	09/03/2012	Nova Laboratories Ltd
XYREM	Sodium oxybate	This product is no longer an orphan medicine. This product was originally an orphan designated on 3 February 2003. Upon request by the MAH, Xyrem has now been removed from the Community Register of Orphan Medicinal products. Cf "List of orphan drugs in Europe with European market authorisation without prior orphan designation in Europe"		

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
YONDELIS	Trabectedin	Treatment of patients with advanced soft tissue sarcoma , after failure of anthracyclines and ifosfamide, or who are unsuited to receive these agents. Efficacy data are based mainly on liposarcoma and leiomyosarcoma patients In combination with pegylated liposomal doxorubicin (PLD), treatment of patients with relapsed platinum-sensitive ovarian cancer	17/09/2007	Pharma Mar S.A.
ZAVESCA	Miglustat	Oral treatment of adult patients with mild to moderate type 1 Gaucher disease . Zavesca may be used only in the treatment of patients for whom enzyme replacement therapy is unsuitable Treatment of progressive neurological manifestations in adult patients and paediatric patients with Niemann-Pick type C disease	20/11/2002	Actelion Registration Ltd

2- Por fecha de la AC en orden descendiente


2012	2009	INCRELEX	2004
BRONCHITOL	CAYSTON	INOVELON	LITAK
DACOGEN	FIRDAPSE (ex-ZENAS)	REVLIMID	LYSODREN
GLYBERA	MEPACT	SIKLOS	PEDEA
JAKAVI	MOZOBIL	SOLIRIS	WILZIN
KALYDECO	NPLATE	TASIGNA	XAGRID
NOVOTHIRTEEN	PEYONA (ex-NYMUSA)	TORISEL	
2008		YONDELIS	2003
REVESTIVE	CEPLENE		ALDURAZYME
SIGNIFOR	FIRAZYR	EVOLTRA	BUSILVEX
XALUPRINE (ex-MER-CAPTOPURINE NOVA)	KUVAN	EXJADE	CARBAGLU
2011	THALIDOMIDE CELGENE	MYOZYME	VENTAVIS
ESBRIET	VIDAZA	NAGLAZYME	
PLENADREN	VOLIBRIS	NEXAVAR	2002
TOBI PODHALER		SAVENE	SOMAVERT
VOTUBIA	2007	SPRYCEL	TRACLEER
VYNDAQEL	ATRIANCE	ORFADIN	TRISENOX
2010	CYSTADANE	PRIALT	ZAVESCA
ARZERRA	DIACOMIT	REVATIO	
TEPADINA	ELAPRASE		2001
VPRIV	GLIOLAN		FABRAZYME



3- Por categoría ATC

A- ALIMENTARY TRACT AND METABOLISM	C- CARDIOVASCULAR SYSTEM	L- ANTOINEPLASTIC AND IMMUNOMODULATING AGENTS	
ALDURAZYME	FIRAZYR	ARZERRA	TEPADINA
CARBAGLU	GLYBERA	ATRIANCE	THALIDOMIDE
CYSTADANE	PEDEA	BUSILVEX	CELGENE
ELAPRASE	TRACLEER	CEPLENE	TORISEL
FABRAZYME	VOLIBRIS	DACOGEN	TRISENOX
KUVAN		ESBRIET	VIDAZA
MYOZYME		EVOLTRA	VOTUBIA
NAGLAZYME	REVATIO	GLIOLAN	XAGRID
ORFADIN		GLIVEC	XALUPRINE (ex-MERCAPTOPURINE NOVA)
REPLAGAL		JAKAVI	YONDELIS
REVESTIVE		LITAK	
VPRIV	INCRELEX	LYSODREN	N- NERVOUS SYSTEM
WILZIN	PLENADREN	MEPACT	DIACOMIT
ZAVESCA	SIGNIFOR	MOZOBIL	FIRDAPSE (ex-ZENAS)
B- BLOOD AND BLOOD FORMING ORGANS	SOMAVERT	NEXAVAR	INOVELON
NOVOTHIRTEEN		REVLIMID	PEYONA (ex-NYMUSA)
NPLATE	CAYSTON	SIKLOS	PRIALT
VENTAVIS	TOBI PODHALER	SOLIRIS	VYNDAQEL
		SPRYCEL	R- RESPIRATORY SYSTEM
		TASIGNA	BRONCHITOL
			KALYDECO
J- GENERAL ANTIINFECTIVES FOR SYSTEMIC USE			V- VARIOUS
			EXJADE
			SAVENE

Number of orphan drugs in Europe with European orphan designation and European marketing authorisation by ATC category



4- Por el titular de la AC

ACTELION REGISTRATION LTD	EPICEPT GMBH	NOVARTIS EUROPHARM LTD	SHIRE ORPHAN THERAPIES GMBH
TRACLEER	CEPLENE	EXJADE	FIRAZYR
ZAVESCA	GENZYME EUROPE B.V.	GLIVEC	SHIRE PHARMACEUTICAL CONTRACTS LTD
ADDMEDICA	ALDURAZYME	JAKAVI	XAGRID
SIKLOS	EVLOTRA	SIGNIFOR	SHIRE PHARMACEUTICALS IRELAND LTD.
ADIENNE S.R.L.	FABRAZYME	TASIGNA	VPRIV
TEPADINA	MOZOBIL	TOBI PODHALER	SPEPHARM HOLDING BV
ALEXION EUROPE SAS	MYOZYME	VOTUBIA	SAVENE
SOLIRIS	GILEAD SCIENCES INTERNATIONAL LIMITED	NOVO NORDISK A/S	SWEDISH ORPHAN BIOVITRUM INTERNATIONAL AB
AMGEN EUROPE B.V.	CAYSTON	NOVOTHIRTEEN	ORFADIN
NPLATE	GLAXO GROUP LTD	NYCOMED DANMARK APS	UNIQUE BIOPHARMA B.V.
BAYER PHARMA AG	ARZERRA	REVESTIVE	GLYBERA
NEXAVAR	ATRIANCE	ORPHAN EUROPE S.A.R.L.	VERTEX PHARMACEUTICALS (U.K.) LIMITED
BAYER SCHERING PHARMA AG	VOLIBRIS	CARBAGLU	KALYDECO
VENTAVIS	IDM PHARMA SAS	CYSTADANE	VIROPHARMA SPRL
BIOCODEX	MEPACT	PEDEA	PLENADREN
DIACOMIT	IPSEN PHARMA	WILZIN	
BIOMARIN EUROPE LTD	INCRELEX	PFIZER LTD	
FIRDAPSE	INTERMUNE UK LTD.	REVATIO	
NAGLAZYME	ESBRIET	SOMAVERT	
BRISTOL-MYERS SQUIBB PHARMA EEWG	JANSSEN-CILAG INTERNATIONAL N V	TORISEL	
SPRYCEL	DACOGEN	PFIZER SPECIALTY UK LTD	
CELGENE EUROPE LTD	LABORATOIRE HRA PHARMA	VYndaqel	
REVLIMID	LYSODREN	PHARMA MAR S.A.	
THALIDOMIDE CELGENE	LIPOMED GMBH	YONDELIS	
VIDAZA	LITAK	PHARMAXIS PHARMACEUTICALS LIMITED	
CEPHALON EUROPE	MEDAC GMBH	BRONCHITOL	
TRISENOX	GLIOLAN	PIERRE FABRE MÉDICAMENT	
CHIESI FARMACEUTICI SPA	MERCK SERONO EUROPE LTD.	BUSILVEX	
PEYONA	KUVAN	SHIRE HUMAN GENETIC THERAPIES AB	
EISAI LTD	NOVA LABORATORIES LTD	ELAPRASE	
INOVELON	XALUPRINE (ex-MER-CAPTOPURINE NOVA)	REPLAGAL	
PRIALT			

Listado de medicamentos huérfanos en Europa con autorización de comercialización europea, sin previa designación huérfana en Europa

1- Por nombre comercial en orden alfabético

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
ADCIRCA	Tadalafil	Treatment of pulmonary arterial hypertension (PAH) classified as WHO functional class II and III, to improve exercise capacity. Efficacy has been shown in idiopathic PAH (IPAH) and in PAH related to collagen vascular disease	30/11/2009	Eli Lilly Nederland B.V.
ADVATE	Octocog alpha	Treatment and prophylaxis of bleeding in patients with haemophilia A (congenital factor VIII deficiency)	02/03/2004	Baxter AG
AFINITOR	Everolimus	Treatment of unresectable or metastatic, well- or moderately-differentiated neuroendocrine tumours of pancreatic origin in adults with progressive disease. Treatment of patients with advanced renal cell carcinoma , whose disease has progressed on or after treatment with VEGF-targeted therapy.	03/08/2009	Novartis Europharm Ltd
ALIMTA	Pemetrexed	In combination with cisplatin for the treatment of chemotherapy naïve patients with unresectable malignant pleural mesothelioma	20/09/2004	Eli Lilly Nederland B.V.
AMMONAPS	Sodium phenylbutyrate	Adjunctive therapy in the chronic management of urea cycle disorders , involving deficiencies of carbamyl phosphate synthetase, ornithine transcarbamylase, or argininosuccinate synthetase. It is indicated in all patients with neonatal-onset presentation (complete enzyme deficiencies, presenting within the first 28 days of life). It is also indicated in patients with late-onset disease (partial enzyme deficiencies, presenting after the first month of life) who have a history of hyperammonaemic encephalopathy	08/12/1999	Swedish Orphan Biovitrum International AB
ATRYN	Antithrombin alpha	Prophylaxis of venous thromboembolism in surgery of patients with congenital antithrombin deficiency , normally given in association with heparin or low molecular weight heparin	28/07/2006	GTC Biotherapeutics UK Limited
AVASTIN	Bevacizumab	In combination with interferon alfa-2a, for first line treatment of patients with advanced and/or metastatic renal cell cancer In combination with carboplatin and paclitaxel, for the front-line treatment of advanced (FIGO stages III B, III C and IV) epithelial ovarian, fallopian tube, or primary peritoneal cancer	12/01/2005	Roche Registration Limited

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
BENEFIX	Recombinant coagulation Factor IX INN = Nonacog alpha	Treatment and prophylaxis of bleeding in patients with haemophilia B (congenital factor IX deficiency)	27/08/1997	Pfizer Ltd
BIOGRASTIM	Filgrastim	In patients, children or adults, with severe congenital, cyclic, or idiopathic neutropenia with an absolute neutrophil count (ANC) of $0.5 \times 10^9/L$, and a history of severe or recurrent infections	15/09/2008	CT Arzneimittel GmbH
BUCCOLAM	Midazolam	Treatment of prolonged, acute, convulsive seizures in infants, toddlers, children and adolescents (from 3 months to < 18 years)	05/09/2011	ViroPharma SPRL
CAELYX	Doxorubicin hydrochloride (pegylated liposomal)	For treatment of advanced ovarian cancer in women who have failed a first-line platinum-based chemotherapy regimen In combination with bortezomib for the treatment of progressive multiple myeloma in patients who have received at least one prior therapy and who have already undergone or are unsuitable for bone marrow transplant Treatment of AIDS-related Kaposi's sarcoma (KS) in patients with low CD4 counts (< 200 CD4 lymphocytes/mm ³) and extensive mucocutaneous or visceral disease	21/06/1996	Janssen-Cilag International N.V.
CANCIDAS	Caspofungin	Treatment of invasive aspergillosis in adult or paediatric patients who are refractory to or intolerant of amphotericin B, lipid formulations of amphotericin B and/or itraconazole. Empirical therapy for presumed fungal infections (such as Candida or Aspergillus) in febrile, neutropaenic adult or paediatric patients	24/10/2001	Merck Sharp & Dohme Ltd
CAPRELSA	Vandetanib	Treatment of thyroid cancer (medullary thyroid cancer)	17/02/2012	AstraZeneca AB
CEPROTIN	Human protein C	In purpura fulminans and coumarin-induced skin necrosis in patients with severe congenital protein C deficiency Short-term prophylaxis in patients with severe congenital protein C deficiency : if surgery or invasive therapy is imminent, while initiating coumarin therapy, when coumarin therapy alone is not sufficient, when coumarin therapy is not feasible	16/07/2001	Baxter AG
CEREZYME	Imiglucerase	Longterm enzyme replacement therapy in patients with a confirmed diagnosis of non-neuronopathic (Type 1) or chronic neuronopathic (Type 3) Gaucher disease and who exhibit clinically significant non-neurological manifestations of the disease, including one or more of the following conditions : anaemia after exclusion of other causes, such as iron deficiency; thrombocytopenia; bone disease after exclusion of other causes such as Vitamin D deficiency; hepatomegaly or splenomegaly	17/11/1997	Genzyme Europe B.V.

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
CINRYZE	C1 inhibitor (human)	Treatment and pre-procedure prevention of angioedema attacks in adults and adolescents with hereditary angioedema (HAE) . Routine prevention of angioedema attacks in adults and adolescents with severe and recurrent attacks of hereditary angioedema (HAE), who are intolerant to or insufficiently protected by oral prevention treatments, or patients who are inadequately managed with repeated acute treatment	15/06/2011	ViroPharma SPRL
COLOBREATHE	Colistimethate sodium	Management of chronic pulmonary infections due to Pseudomonas aeruginosa in patients with cystic fibrosis (CF) aged 6 years and older	13/02/2012	Forest Laboratories UK Ltd
CYSTAGON	Mercaptamine bitartrate	Treatment of proven nephropathic cystinosis Cysteamine reduces cystine accumulation in some cells (e.g. leukocytes, muscle and liver cells) of nephropathic cystinosis patients and, when treatment is started early, it delays the development of renal failure	23/06/1997	Orphan Europe S.A.R.L.
DUKORAL	Vibrio cholerae and recombinant cholera toxin B-subunit	Active immunisation against disease caused by Vibrio cholerae serogroup O1 in adults and children from 2 years of age who will be visiting endemic/epidemic areas	28/04/2004	Crucell Sweden AB
ENBREL	Etanercept	Treatment of polyarthritis (rheumatoid-factor-positive or -negative) and extended oligoarthritis in children and adolescents from the age of 2 years who have had an inadequate response to, or who have proved intolerant of, methotrexate Treatment of psoriatic arthritis in adolescents from the age of 12 years who have had an inadequate response to, or who have proved intolerant of, methotrexate Treatment of enthesitis-related arthritis in adolescents from the age of 12 years who have had an inadequate response to, or who have proved intolerant of, conventional therapy. Enbrel has not been studied in children aged less than 2 years	03/02/2000	Pfizer Ltd
ERBITUX	Cetuximab	Treatment of patients with squamous cell cancer of the head and neck , in combination with radiation therapy for locally advanced disease and in combination with platinum-based chemotherapy for recurrent and/or metastatic disease	29/06/2004	Merck KGaA
EURARTESIM	Piperaquine tetraphosphate / dihydroartemisinin	Treatment of uncomplicated Plasmodium falciparum malaria in adults, children and infants 6 months and over and weighing 5 kg or more.	27/10/2011	Sigma-Tau Industrie Farmaceutiche Riunite S.p.A
FERRIPROX	Deferiprone	Treatment of iron overload in patients with thalassaemia major when deferoxamine therapy is contraindicated or inadequate	25/08/1999	Apotex Europe B.V.

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
FILGRASTIM HEXAL	Filgrastim	In patients, children or adults, with severe congenital, cyclic, or idiopathic neutropenia with an absolute neutrophil count (ANC) of $0.5 \times 10^9/l$, and a history of severe or recurrent infections	06/02/2009	Hexal AG
FLEBOGAMMA DIF	Human normal immunoglobulin	Replacement therapy in: Primary immunodeficiency syndromes such as: - congenital agammaglobulinaemia and hypogammaglobulinaemia - common variable immunodeficiency - severe combined immunodeficiency - Wiskott Aldrich syndrome Myeloma or chronic lymphocytic leukaemia with severe secondary hypogammaglobulinaemia and recurrent infections. Immunomodulation in: Idiopathic thrombocytopenic purpura (ITP), in children or adults at high risk of bleeding or prior to surgery to correct the platelet count. Guillain Barré syndrome Kawasaki disease	23/07/2007	Instituto Grifols S.A.
GONAL-F	Recombinant human follicle stimulating hormone INN = Follitropin alpha	Stimulation of spermatogenesis in men who have congenital or acquired hypogonadotropic hypogonadism with concomitant human Chorionic Gonadotrophin (hCG) therapy	20/10/1995	Merck Serono Europe Ltd
HELIXATE NEXGEN	Octocog alpha	Treatment and prophylaxis of bleeding in patients with haemophilia A (congenital factor VIII deficiency)	04/08/2000	Bayer Pharma AG
HERCEPTIN	Trastuzumab	In combination with capecitabine or 5-fluorouracil and cisplatin is indicated for the treatment of patients with HER2 positive metastatic adenocarcinoma of the stomach or gastro-esophageal junction who have not received prior anti-cancer treatment for their metastatic disease. Herceptin should only be used in patients with metastatic gastric cancer whose tumours have HER2 overexpression as defined by IHC2+ and a confirmatory SISH or FISH result, or by an IHC 3+ result. Accurate and validated assay methods should be used.	28/08/2000	Roche Registration Limited
HIZENTRA	Human normal immunoglobulin (SCIg)	Replacement therapy in adults and children in primary immunodeficiency syndromes such as: - congenital agammaglobulinaemia and hypogammaglobulinaemia - common variable immunodeficiency - severe combined immunodeficiency - IgG subclass deficiencies with recurrent infections - Replacement therapy in myeloma or chronic lymphocytic leukaemia with severe secondary hypogammaglobulinaemia and recurrent infections	14/04/2011	CSL Behring GmbH

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
HUMIRA	Adalimumab	In combination with methotrexate is indicated for the treatment of active polyarticular juvenile idiopathic arthritis , in children and adolescents aged 4 to 17 years who have had an inadequate response to one or more disease-modifying antirheumatic drugs (DMARDs) As monotherapy in case of intolerance to methotrexate or when continued treatment with methotrexate is inappropriate Humira has not been studied in children aged less than 4 years	08/09/2003	Abbott Laboratories Ltd
HYCAMTIN	Topotecan	Treatment of patients with metastatic carcinoma of the ovary after failure of first-line or subsequent therapy Treatment of patients with relapsed small cell lung cancer [SCLC] for whom re-treatment with the first-line regimen is not considered appropriate In combination with cisplatin for patients with carcinoma of the cervix recurrent after radiotherapy and for patients with Stage IVB disease. Patients with prior exposure to cisplatin require a sustained treatment free interval to justify treatment with the combination HYCAMTIN capsules are indicated as monotherapy for the treatment of adult patients with relapsed small cell lung cancer (SCLC) for whom re-treatment with the first-line regimen is not considered appropriate	12/11/1996	SmithKline Beecham Ltd
ILARIS	Canakinumab	Treatment of Cryopyrin-Associated Periodic Syndromes (CAPS) in adults, adolescents and children aged 4 years and older with body weight above 15 kg, including: - Muckle-Wells Syndrome (MWS), - Neonatal-Onset Multisystem Inflammatory Disease (NOMID) / Chronic Infantile Neurological, Cutaneous, Articular Syndrome (CINCA), - Severe forms of Familial Cold Autoinflammatory Syndrome (FCAS) / Familial Cold Urticaria (FCU) presenting with signs and symptoms beyond cold-induced urticarial skin rash	23/10/2009	Novartis Europarm Ltd
INLYTA	Axitinib	For the treatment of adult patients with advanced renal cell carcinoma (RCC) after failure of prior treatment with sunitinib or a cytokine	03/09/2012	Pfizer Ltd.

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
INOMAX	Nitric oxide	In conjunction with ventilatory support and other appropriate active substances: - for the treatment of newborn infants ≥34 weeks gestation with hypoxic respiratory failure associated with clinical or echocardiographic evidence of pulmonary hypertension , in order to improve oxygenation and to reduce the need for extracorporeal membrane oxygenation. - as part of the treatment of peri- and post-operative pulmonary hypertension in adults and newborn infants, infants and toddlers, children and adolescents, ages 0-17 years in conjunction to heart surgery, in order to selectively decrease pulmonary arterial pressure and improve right ventricular function and oxygenation	01/08/2001	INO Therapeutics AB
INTRONA	Interferon alpha-2b	Treatment of patients with hairy cell leukaemia Monotherapy treatment of adults with Philadelphia chromosome or bcr/abl translocation positive chronic myelogenous leukaemia Combination therapy with cytarabine administered during the first 12 months of treatment has been demonstrated to significantly increase the rate of major cytogenetic responses and to significantly prolong the overall survival at three years when compared to interferon alfa-2b monotherapy Treatment of patients with multiple myeloma , as maintenance therapy in patients who have achieved objective remission (more than 50 % reduction in myeloma protein) following initial induction chemotherapy Treatment of high tumour burden follicular lymphoma as adjunct to appropriate combination induction chemotherapy such as a CHOP-like regimen Treatment of carcinoid tumours with lymph node or liver metastases and with "carcinoid syndrome"	09/03/2000	Merck Sharp & Dohme Limited
IXIARO	Japanese Encephalitis Vaccine (inactivated, adsorbed)	For active immunization against Japanese encephalitis for adults	31/03/2009	Intercell AG
KEPPRA	Levetiracetam	As monotherapy in the treatment of partial onset seizures with or without secondary generalisation in patients from 16 years of age with newly diagnosed epilepsy . As adjunctive therapy in the treatment of partial onset seizures with or without secondary generalisation in adults, children and infants from 1 month of age with epilepsy ; in the treatment of myoclonic seizures in adults and adolescents from 12 years of age with Juvenile Myoclonic Epilepsy ; in the treatment of primary generalised tonic-clonic seizures in adults and adolescents from 12 years of age with Idiopathic Generalised Epilepsy	29/09/2000	UCB Pharma SA

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
KIOVIG	Human normal immunoglobulin	<p>Replacement therapy in adults, and children and adolescents (0-18 years) in:</p> <ul style="list-style-type: none"> - Primary immunodeficiency syndromes with impaired antibody production - Hypogammaglobulinaemia and recurrent bacterial infections in patients with chronic lymphocytic leukaemia, in whom prophylactic antibiotics have failed. - Hypogammaglobulinaemia and recurrent bacterial infections in plateau phase multiple myeloma patients who have failed to respond to pneumococcal immunisation. - Hypogammaglobulinaemia in patients after allogeneic haematopoietic stem cell transplantation (HSCT). <p>Immunomodulation in adults, and children and adolescents (0-18 years) in:</p> <ul style="list-style-type: none"> - Primary immune thrombocytopenia (ITP), in patients at high risk of bleeding or prior to surgery to correct the platelet count. - Guillain Barré syndrome. - Kawasaki disease. - Multifocal Motor Neuropathy (MMN). 	19/01/2006	Baxter AG
KOGENATE BAYER	Octocog alpha	Treatment and prophylaxis of bleeding in patients with haemophilia A (congenital factor VIII deficiency)	04/08/2000	Bayer Pharma AG
MABTHERA	Rituximab	<p>Indicated in adults for:</p> <p>Treatment of previously untreated patients with stage III-IV follicular lymphoma in combination with chemotherapy</p> <p>Maintenance therapy is indicated for the treatment of follicular lymphoma patients responding to induction therapy</p> <p>Monotherapy is indicated for treatment of patients with stage III-IV follicular lymphoma who are chemoresistant or are in their second or subsequent relapse after chemotherapy</p> <p>Treatment of patients with CD20 positive diffuse large B cell non-Hodgkin's lymphoma in combination with CHOP (cyclophosphamide, doxorubicin, vincristine, prednisolone) chemotherapy</p> <p>In combination with chemotherapy, treatment of patients with previously untreated and relapsed/refractory chronic lymphocytic leukaemia (CLL).</p> <p>Only limited data are available on efficacy and safety for patients previously treated with monoclonal antibodies including MabThera or patients refractory to previous MabThera plus chemotherapy</p>	02/06/1998	Roche Registration Limited

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
NIVESTIM	Filgrastim	In patients, children or adults, with severe congenital, cyclic, or idiopathic neutropenia with an absolute neutrophil count (ANC) of $0.5 \times 10^9/l$, and a history of severe or recurrent infections	08/06/2010	Hospira UK Ltd
NOVOSEVEN	Human recombinant coagulation Factor VIIa INN = Eptacog alpha (activated)	Treatment of bleeding episodes and for the prevention of bleeding in those undergoing surgery or invasive procedures in the following patient groups : in patients with congenital haemophilia with inhibitors to coagulation factors VIII or IX > 5 BU; in patients with congenital haemophilia who are expected to have a high anamnestic response to factor VIII or factor IX administration; in patients with acquired haemophilia ; in patients with congenital FVII deficiency ; in patients with Glanzmann's thrombasthenia with antibodies to GP IIb - IIIa and/or HLA, and with past or present refractoriness to platelet transfusions	23/02/1996	Novo Nordisk A/S
NOXAFIL	Posaconazole	Treatment of the fungal infections in adults: <ul style="list-style-type: none"> - Invasive aspergillosis in patients with disease that is refractory to amphotericin B or itraconazole or in patients who are intolerant of these medicinal products - Fusariosis in patients with disease that is refractory to amphotericin B or in patients who are intolerant of amphotericin B - Chromoblastomycosis and mycetoma in patients with disease that is refractory to itraconazole or in patients who are intolerant of itraconazole - Coccidioidomycosis in patients with disease that is refractory to amphotericin B, itraconazole or fluconazole or in patients who are intolerant of these medicinal products Prophylaxis of invasive fungal infections in : <ul style="list-style-type: none"> - Patients receiving remission-induction chemotherapy for acute myelogenous leukemia (AML) or myelodysplastic syndromes (MDS) expected to result in prolonged neutropenia and who are at high risk of developing invasive fungal infections - Hematopoietic stem cell transplant (HSCT) recipients who are undergoing high-dose immunosuppressive therapy for graft versus host disease and who are at high risk of developing invasive fungal infections 	25/10/2005	Merck Sharp & Dohme Ltd.
OMNITROPE	Somatropin	Growth disturbance due to insufficient secretion of growth hormone (GH) and growth disturbance associated with Turner syndrome or chronic renal insufficiency. Prader-Willi syndrome (PWS), for improvement of growth and body composition. Replacement therapy in adults with pronounced growth hormone deficiency (patients with known hypothalamic pituitary pathology and at least one known deficiency of a pituitary hormone not being prolactin)	12/04/2006	Sandoz GmbH

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
ORENCIA	Abatacept	In combination with methotrexate, for the treatment of moderate to severe active polyarticular juvenile idiopathic arthritis (JIA) in paediatric patients 6 years of age and older who have had an insufficient response to other DMARDs including at least one TNF inhibitor	21/05/2007	Bristol-Myers Squibb Pharma EEIG
OZURDEX	Dexamethasone	For the treatment of adult patients with inflammation of the posterior segment of the eye presenting as non-infectious uveitis	27/07/2010	Allergan Pharmaceuticals Ireland
PANRETIN	Alitretinoin	Topical treatment of cutaneous lesions in patients with AIDS-related Kaposi's sarcoma (KS) : when lesions are not ulcerated or lymphoedematous, and treatment of visceral KS is not required, and when lesions are not responding to systemic antiretroviral therapy, and radiotherapy or chemotherapy are not appropriate	11/10/2000	Eisai Ltd
PIXUVRI	Pixantrone dimaleate	As monotherapy for the treatment of adult patients with multiply relapsed or refractory aggressive non-Hodgkin B cell lymphomas (NHL). The benefit of pixantrone treatment has not been established in patients when used as fifth line or greater chemotherapy in patients who are refractory to last therapy	10/05/2012	CTI Life Sciences Ltd
PRIVIGEN	Human normal immunoglobulin (IVIg)	Replacement therapy in : <ul style="list-style-type: none"> - Primary immunodeficiency (PID) syndromes such as: <ul style="list-style-type: none"> - congenital agammaglobulinaemia and hypogammaglobulinaemia - common variable immunodeficiency - severe combined immunodeficiency - Wiskott Aldrich syndrome - Myeloma or chronic lymphocytic leukaemia with severe secondary hypogammaglobulinaemia and recurrent infections. Immunomodulation in : <ul style="list-style-type: none"> - Immune thrombocytopenic purpura (ITP), in children or adults at high risk of bleeding or prior to surgery to correct the platelet count - Guillain-Barré syndrome - Kawasaki disease 	25/04/2008	CSL Behring GmbH
PUREGON	Follitropin beta	Treatment of deficient spermatogenesis due to hypogonadotropic hypogonadism	03/05/1996	NV Organon
RATIOGRASTIM	Filgrastim	In patients, children or adults, with severe congenital, cyclic, or idiopathic neutropenia with an absolute neutrophil count (ANC) of $0.5 \times 10^9/l$, and a history of severe or recurrent infections	15/09/2008	Ratiopharm GmbH
REFACTO AF	Moroctocog alpha	Treatment and prophylaxis of bleeding in patients with haemophilia A (congenital factor VIII deficiency) in adults and children of all ages, including newborns	13/04/1999	Pfizer Ltd

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
REVOLADE	Eltrombopag	For adult chronic immune (idiopathic) thrombocytopenic purpura (ITP) splenectomised patients who are refractory to other treatments (e.g. corticosteroids, immunoglobulins). Revolade may be considered as second line treatment for adult non-splenectomised patients where surgery is contraindicated This product is no longer an orphan medicine.	11/03/2010	Glaxo-SmithKline Trading Services Limited
RILUTEK	Riluzole	To extend life or the time to mechanical ventilation for patients with amyotrophic lateral sclerosis (ALS)	10/06/1996	Aventis Pharma S.A.
ROACTEMRA	Tocilizumab	Treatment of active systemic juvenile idiopathic arthritis (sJIA) in patients 2 years of age and older, who have responded inadequately to previous therapy with NSAIDs and systemic corticosteroids. RoActemra can be given as monotherapy (in case of intolerance to MTX or where treatment with MTX is inappropriate) or in combination with MTX.	16/01/2009	Roche Registration Ltd
RUCONEST	Conestat alfa	Treatment of acute angioedema attacks in adults with hereditary angioedema (HAE) due to C1 esterase inhibitor deficiency	28/10/2010	Pharming Group N.V.
SAMSCA	Tolvaptan	Treatment of adult patients with hyponatraemia secondary to syndrome of inappropriate antidiuretic hormone secretion (SIADH)	03/08/2009	Otsuka Pharmaceutical Europe Ltd
SUTENT	Sunitinib	Treatment of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST) after failure of imatinib mesilate treatment due to resistance or intolerance Treatment of advanced/metastatic renal cell carcinoma (MRCC) in adults Treatment of unresectable or metastatic, well-differentiated pancreatic neuroendocrine tumours (pNET) with disease progression in adults Experience with SUTENT as first-line treatment is limited	19/07/2006	Pfizer Limited
TARCEVA	Erlotinib	In combination with gemcitabine, for the treatment of patients with metastatic pancreatic cancer . When prescribing Tarceva, factors associated with prolonged survival should be taken into account . No survival advantage could be shown for patients with locally advanced disease	19/09/2005	Roche Registration Limited
TARGRETIN	Bexarotene	Treatment of skin manifestations of advanced stage cutaneous T-cell lymphoma (CTCL) patients refractory to at least one systemic treatment	29/03/2001	Eisai Ltd
TAXOTERE	Docetaxel	In combination with cisplatin and 5-fluorouracil for the treatment of patients with metastatic gastric adenocarcinoma , including adenocarcinoma of the gastroesophageal junction, who have not received prior chemotherapy for metastatic disease In combination with cisplatin and 5-fluorouracil for the induction treatment of patients with locally advanced squamous cell carcinoma of the head and neck	27/11/1995	Aventis Pharma S.A.


TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
TEMODAL	Temozolomide	Treatment of adult patients with newly-diagnosed glioblastoma multiforme concomitantly with radiotherapy (RT) and subsequently as monotherapy treatment Treatment of children from the age of three years, adolescents and adult patients with malignant glioma , such as glioblastoma multiforme or anaplastic astrocytoma , showing recurrence or progression after standard therapy	26/01/1999	Schering-Plough Europe
TEVAGRASTIM	Filgrastim	In patients, children or adults, with severe congenital, cyclic, or idiopathic neutropenia with an absolute neutrophil count (ANC) of $0.5 \times 10^9/L$, and a history of severe or recurrent infections	15/09/2008	Teva Generics GmbH
TEYSUNO	Tegafur/Gimeracil /Oteracil	In adults for the treatment of advanced gastric cancer when given in combination with cisplatin	14/03/2011	Nordic Group BV
THYROGEN	Thyrotropin alfa	For use with serum thyroglobulin (Tg) testing with or without radioiodine imaging for the detection of thyroid remnants and well-differentiated thyroid cancer in post-thyroidectomy patients maintained on hormone suppression therapy (THST). Low-risk patients with well-differentiated thyroid carcinoma who have undetectable serum Tg levels on THST and no rh (recombinant human) TSH-stimulated increase of Tg levels may be followed-up by assaying rh TSH-stimulated Tg levels. For pre-therapeutic stimulation in combination with a range of 30 mCi (1.1 GBq) to 100 mCi (3.7 GBq) radioiodine for ablation of thyroid tissue remnants in patients who have undergone a near-total or total thyroidectomy for well-differentiated thyroid cancer and who do not have evidence of distant metastatic thyroid cancer	09/03/2000	Genzyme Europe B.V.
VALTROPIN	Somatropin	Long-term treatment of children with growth failure due to an inadequate secretion of normal endogenous growth hormone Treatment of short stature in children with Turner syndrome , confirmed by chromosome analysis Replacement therapy in adults with pronounced growth hormone deficiency of either childhood- or adult-onset aetiology (patients with known hypothalamic-pituitary pathology and at least one additional known deficiency of a pituitary hormone not being prolactin)	24/04/2006	BioPartners GmbH
VEDROP	Tocoferolsolan	Indicated in vitamin E deficiency due to digestive malabsorption in paediatric patients suffering from congenital chronic cholestasis or hereditary chronic cholestasis , from birth (in term newborns) to 16 or 18 years of age, depending on the region	24/07/2009	Orphan Europe S.A.R.L

TRADENAME	ACTIVE SUBSTANCE	MARKETING AUTHORISATION INDICATION	MARKETING AUTHORISATION DATE (DD/MM/YYYY)	MARKETING AUTHORISATION HOLDER
VELCADE	Bortezomib	In combination with melphalan and prednisone for the treatment of patients with previously untreated multiple myeloma who are not eligible for high-dose chemotherapy with bone marrow transplant As mono-therapy for the treatment of progressive multiple myeloma in patients who have received at least 1 prior therapy and who have already undergone or are unsuitable for bone marrow transplantation	26/04/2004	Janssen-Cilag International NV
VFEND	Voriconazole	For treatment of invasive aspergillosis For treatment of serious fungal infections caused by <i>Scedosporium</i> spp. and <i>Fusarium</i> spp. (Fusariosis) VFEND should be administered primarily to patients with progressive, possibly life-threatening infections	19/03/2002	Pfizer Limited
VOTRIENT	Pazopanib	In adults for the first-line treatment of advanced renal cell carcinoma (RCC) and for patients who have received prior cytokine therapy for advanced disease. For the treatment of adult patients with selective subtypes of advanced soft-tissue sarcoma (STS) who have received prior chemotherapy for metastatic disease or who have progressed within 12 months after (neo)-adjuvant therapy. Efficacy and safety have only been established in certain STS histological tumour subtypes	14/06/2010	Glaxo Group Ltd
XELODA	Capecitabine	First-line treatment of advanced gastric cancer in combination with a platinum-based regimen	02/02/2001	Roche Registration Limited
XYREM	Sodium oxybate	Treatment of narcolepsy with cataplexy in adult patients	13/10/2005	UCB Pharma Ltd
ZARZIO	Filgrastim	In patients, children or adults, with severe congenital, cyclic, or idiopathic neutropenia with an absolute neutrophil count (ANC) of $0.5 \times 10^9/l$, and a history of severe or recurrent infections	06/02/2009	Sandoz GmbH
ZEVALIN	Ibritumomab tiuxetan	Consolidation therapy after remission induction in previously untreated patients with follicular lymphoma Treatment of adult patients with rituximab relapsed or refractory CD20+ follicular B-cell non-Hodgkin's lymphoma (NHL)	16/01/2004	Bayer Pharma AG
ZUTECTRA	Human Hepatitis B Immunoglobulin	Prevention of hepatitis B virus (HBV) re-infection in HBV-DNA negative patients over 6 months after liver transplantation for hepatitis B induced liver failure . Zutectra is indicated in adults only. The concomitant use of adequate virostatic agents should be considered, if appropriate, as standard of hepatitis B re-infection prophylaxis	30/11/2009	Biotest Pharma GmbH

2- Por fecha de la AC en orden descendiente

2012	ROACTEMRA	2004	PANRETIN
CAPRELSA	SAMSCA	ADVATE	THYROGEN
COLOBREATHE	VEDROP	ALIMTA	1999
INLYTA	ZARZIO	DUKORAL	AMMONAPS
PIXUVRI	ZUTECTRA	ERBITUX	FERRIPROX
2011	2008	VELCADE	REFACTO AF
BUCCOLAM	BIOGRASTIM	ZEVALIN	TEMODAL
CINRYZE	PRIVIGEN	HUMIRA	1998
EURARTESIM	RATIOGRASTIM	VFEND	MABTHERA
HIZENTRA	TEVAGRASTIM	2001	1997
TEYSUNO	2007	CANCIDAS	BENEFIX
2010	FLEBOGAMMA DIF	CEPROTIN	CEREZYME
NIVESTIM	ORENCIA	INOMAX	CYSTAGON
OZURDEX	2006	TARGRETIN	1996
REVOLADE	ATRYN	XELODA	CAELYX
RUCONEST	KIOVIG	2000	HYCAMTIN
VOTRIENT	OMNITROPE	ENBREL	NOVOSEVEN
2009	SUTENT	HELIXATE NEXGEN	PUREGON
ADCIRCA	VALTROPIN	HERCEPTIN	RILUTEK
AFINITOR	2005	INTRONA	1995
FILGRASTIM HEXAL	AVASTIN	KEPPRA	GONAL-F
ILARIS	NOXAFL	KOGENATE BAYER	TAXOTERE
IXIARO	TARCEVA		
	XYREM		


Number of orphan drugs in Europe with European market authorisation without prior orphan designation in Europe by date of MA



3- Por categoría ATC

A- ALIMENTARY TRACT AND METABOLISM	H- SYSTEMIC HORMONAL PREPARATIONS, EXCL, SEX HORMONES AND INSULINS		N- NERVOUS SYSTEM
AMMONAPS	OMNITROPE	ENBREL	BUCCOLAM
CEREZYME	THYROGEN	ERBITUX	KEPPRA
CYSTAGON	VALTROPIN	FILGRASTIM HEXAL	RILUTEK
VEDROP	J- GENERAL ANTIINFECTIVES FOR SYSTEMIC USE	HERCEPTIN	XYREM
B- BLOOD AND BLOOD FORMING ORGANS	CANCIDAS	HUMIRA	P- ANTIPARASITIC PRODUCTS, INSECTICIDES AND REPELLENTS
ADVATE	DUKORAL	HYCAMTIN	EURARTESIM
ATRYN	FLEBOGAMMA DIF	ILARIS	R- RESPIRATORY SYSTEM
BENEFIX	HIZENTRA	INLYTA	COLOBREATHE
CEPROTIN	IXIARO	INTRONA	INOMAX
CINRYZE	KIOVIG	MABTHERA	S- SENSORY ORGANS
HELIXATE NEXGEN	NOXAFILE	NIVESTIM	OZURDEX
KOGENATE BAYER	PRIVIGEN	ORENCIA	V- VARIOUS
NOVOSEVEN	VFEND	PANRETIN	FERRIPROX
REFACTO AF	ZUTECTRA	PIXUVRI	ZEVALIN
REVOLADE	L- ANTOINEPLASTIC AND IMMUNOMODULATING AGENTS	RATIOGRASTIM	
RUCONEST	AFINITOR	ROACTEMRA	
C- CARDIOVASCULAR SYSTEM	ALIMTA	SUTENT	
SAMSCA	AVASTIN	TARCEVA	
G- GENITO URINARY SYSTEM AND SEX HORMONES	BIOGRASTIM	TARGRETIN	
ADCIRCA	CAELYX	TAXOTERE	
GONAL-F	CAPRELSA	TEMODAL	
PUREGON		TEVAGRASTIM	
		TEYSUNO	
		VELCADE	
		VOTRIENT	
		XELODA	
		ZARZIO	

Number of orphan drugs in Europe with European market authorisation without prior orphan designation in Europe by ATC category



4- Por el titular de la AC

ABBOTT LABORATORIES LTD	EISAI LTD	MERCK KGAA	RATIOPHARM GMBH
HUMIRA	PANRETIN	ERBITUX	RATIOGRASTIM
ALLERGAN PHARMACEUTICALS IRELAND	TARGETIN	MERCK SERONO EUROPE LTD	ROCHE REGISTRATION LTD
OZURDEX	ELI LILLY NEDERLAND B.V.	GONAL-F	AVASTIN
APOTEX EUROPE B.V.	ADCIRCA	MERCK SHARP & DOHME LTD	HERCEPTIN
FERRIPROX	ALIMTA	CANCIDAS	MABTHERA
ASTRAZENECA AB	FOREST LABORATORIES UK LTD	INTRONA	ROACTEMRA
CAPRELSA	COLOBREATHE	NOXAFIL	TARCEVA
AVENTIS PHARMA S.A.	GENZYME EUROPE B.V.	NORDIC GROUP BV	XELODA
RILUTEK	CEREZYME	TEYSUNO	SANDOZ GMBH
TAXOTERE	THYROGEN	NOVARTIS EUROPHARM LTD	OMNITROPE
BAXTER AG	GLAXO GROUP LTD	AFINITOR	ZARZIO
ADVATE	VOTRIENT	ILARIS	SCHERING-PLOUGH EUROPE
CEPROTIN	GLAXOSMITHKLINE TRADING SERVICES LIMITED	NOVO NORDISK A/S	TEMODAL
KIOVIG	REVOLADE	NOVOSEVEN	SMITHKLINE BEECHAM LTD
BAYER PHARMA AG	GTC BIOTHERAPEUTICS UK LIMITED	NV ORGANON	HYCAMTIN
HELIXATE NEXGEN	ATRYN	PUREGON	SIGMA-TAU INDUSTRIE FARMACEUTICHE RIUNITE S.P.A
KOGENATE BAYER	HEXAL AG	ORPHAN EUROPE S.A.R.L.	EURARTESIM
ZEVALIN	FILGRASTIM HEXAL	CYSTAGON	SWEDISH ORPHAN BIOVITRUM INTERNATIONAL AB
BIOPARTNERS GMBH	HOSPIRA UK LTD	VEDROP	AMMONAPS
VALTROPIN	NIVESTIM	OTSUKA PHARMACEUTICAL EUROPE LTD	TEVA GENERICS GMBH
BIOTEST PHARMA GMBH	INO THERAPEUTICS AB	SAMSCA	TEVAGRASTIM
ZUTECTRA	INOMAX	PFIZER LTD	UCB PHARMA LTD
BRISTOL-MYERS SQUIBB PHARMA EEWG	INSTITUTO GRIFOLS S.A.	BENEFIX	XYREM
ORENCIA	FLEBOGAMMA DIF	ENBREL	UCB PHARMA SA
CRUCELL SWEDEN AB	INTERCELL AG	INLYTA	KEPPRA
DUKORAL	IXIARO	REFACTO AF	VIROPHARMA SPRL
CSL BEHRING GMBH	JANSSEN-CILAG INTERNATIONAL NV	SUTENT	BUCCOLAM
HIZENTRA	CAELYX	VFEND	CINRYZE
PRIVIGEN	VELCADE	PHARMING GROUP N.V.	
CT ARZNEIMITTEL GMBH		RUCONEST	
BIOGRASTIM			
CTI LIFE SCIENCES LTD			
PIXUVRI			

Para cualquier pregunta o comentario, por favor contacte con: contact.orphanet@inserm.fr

Redactora jefe : Odile Kremp ● Redactora : Virginie Hivert ● Diseño : Céline Angin ● Fotografía : M. Depardieu/Inserm

La forma adecuada para citar este documento es la siguiente :

« Listados de Medicamentos Huérfanos en Europa », Informes Periódicos de Orphanet, Serie *Medicamentos Huérfanos*, Octubre 2012,
http://www.orpha.net/orphacom/cahiers/docs/ES/listado_de_medicamentos_huerfanos_in_Europa.pdf